

## Histone H4 Biotin Conjugated Peptide - 000-006-K86

**Code:** 000-006-K86

**Size:** 1 mg

**Product Description:** Histone H4 Biotin Conjugated Peptide - 000-006-K86

**Concentration:** 1.0 mg/mL by dry weight

**PhysicalState:** Lyophilized

<b>Label</b>	Biotin
<b>Buffer</b>	None
<b>Reconstitution Volume</b>	1.0 mL
<b>Reconstitution Buffer</b>	Restore with deionized water (or equivalent)
<b>Storage Condition</b>	Store vial at 2 - 8 ° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Dilute only prior to immediate use.
<b>Synonyms</b>	H4/A, HIST2H4, HIST2H4A, HIST1H4A, histone cluster 4, H4, histone 4, Histone H4 peptide, histone H4, MGC24116, control peptide, blocking peptide
<b>Application Note</b>	Histone H4 Control Peptide is suitable for use in ELISA, Western Blot, Dot blot, PCA, and other assays. Control peptide should be used at 1.0 µg per 1.0 µl of antiserum in per assay. Specific conditions for reactivity should be optimized by the end user.
<b>Background</b>	The nucleosome is comprised of 146 bp of DNA wrapped around a series of histone proteins arranged as an octamer consisting of 2 copies of histone H2A, H2B, H3 and H4. Within the nucleosome core the histone proteins are covalent modified at specific residues predominantly within the N-terminal tail including lysine (acetylation, methylation, SUMOylation, and ubiquitinylation), arginine methylation and citrullination, serine and threonine phosphorylation, as well as proline isomerization. The lysine side chains can carry up to three methyl groups (mono-, di- and tri- methylated forms) and the arginine side chain can be monomethylated or can be dimethylated as the symmetric or asymmetric forms. The modifications show temporal, disease-specific, and other types of cell-specific regulation and there are specific families of enzymes that regulate the methylation, demethylation, acetylation, deacetylation and other modifications. Research has indicated that whereas the histone mark H3K4Me3 (tri-methyl lysine 4 of histone H3) localizes to gene promoter regions (it is associated with transcriptional activation) other modifications at H3K4 such as monomethyl is present predominantly at enhancer sequences. Specific marks have been associated with activation (H3K9Me1, H3K27Me1, and H4K20Me1) or repression (H3K9Me2 and Me3, H3K27Me2 and Me3, and H4K20Me2 and Me3) of genes. Monomethylation of H4 at K20, catalyzed by SET8, is essential to genome replication and stability. Multiple DNA breaks are associated with demethylation at this site, resulting in activation of p53 to avoid mitosis and aberrant chromosomal activity. In mammalian stem cells, Xist expression blocks the formation of H4K20me1, which is one of the first examples of a direct connection between chromatin and stem cell differentiation. Histone H4 are ideal for researchers interested in Chromatin Research, DNA replication Transcription Translation and Splicing, Epigenetics, Chromatin Modifiers, Histones and Modified Histones, DNA Repair, and HA Epitope Tags research.
<b>Purity And Specificity</b>	Greater than 95% specific peptide.
<b>Assay Dilutions</b>	Control peptide should be used at 1.0 µg per 1.0 µl of antiserum per assay.
<b>Other Assays</b>	Control peptide should be used at 1.0 µg per 1.0 µl of antiserum per assay.
<b>Expiration</b>	Expiration date is one (1) year from date of opening.
<b>Related Products</b>	
611-1302	Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302
BSA-30	BOVINE SERUM ALBUMIN 30% Solution - BSA-30
MB-070	Blocking Buffer for Fluorescent Western Blotting - MB-070
100-401-E81	Histone H3 Antibody100-401-E81

**Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.