



## CAT IgG F(c) fragment Peroxidase conjugated - 002-0303

**Code:** 002-0303

**Size:** 1 mg

**Product Description:** CAT IgG F(c) fragment Peroxidase conjugated - 002-0303

**Concentration:** 1.0 mg/mL by UV absorbance at 280 nm

**PhysicalState:** Lyophilized

|                               |                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|-------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Label</b>                  | Peroxidase (Horseradish)                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>Buffer</b>                 | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2                                                                                                                                                                                                                                                                                                                                                                                                              |
| <b>Reconstitution Volume</b>  | 1.0 mL                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Reconstitution Buffer</b>  | Restore with deionized water (or equivalent)                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Stabilizer</b>             | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Preservative</b>           | 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!                                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>Storage Condition</b>      | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.                                                                                                                         |
| <b>Synonyms</b>               | Cat IgG F(c) fragment HRP conjugated, Cat IgG F(c) fragment peroxidase conjugated                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Application Note</b>       | Cat IgG F(c) fragment Peroxidase conjugated can be utilized as a control or standard reagent in Western Blotting and ELISA experiments.                                                                                                                                                                                                                                                                                                                                 |
| <b>Background</b>             | Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. This antibody possesses the F(c) region, recognized by high-affinity Fc receptor proteins. |
| <b>Purity And Specificity</b> | This product was prepared from normal serum by delipidation, salt fractionation, ion exchange chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Cat IgG, anti-Cat IgG F(c) and anti-Cat Serum. No reaction was observed against anti-Cat IgG F(ab') <sub>2</sub> or anti-Papain.                                     |
| <b>Assay Dilutions</b>        | User Optimized                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Other Assays</b>           | User Optimized                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Expiration</b>             | Expiration date is one (1) year from date of opening.                                                                                                                                                                                                                                                                                                                                                                                                                   |

### Related Products

|          |                                                                               |
|----------|-------------------------------------------------------------------------------|
| 010-0102 | MOUSE IgG whole molecule - 010-0102                                           |
| 610-4302 | Anti-MOUSE IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated - 610-4302       |
| 611-1302 | Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302        |
| BSA-50   | BOVINE SERUM ALBUMIN - Fraction V (Immunoglobulin and Protease Free) - BSA-50 |

### Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.