

DOG IgG F(c) fragment - 004-0103
Code: 004-0103

Size: 1 mg

Product Description: DOG IgG F(c) fragment - 004-0103

Concentration: 1.2 mg/mL by UV absorbance at 280 nm

PhysicalState: Liquid (sterile filtered)

Label	Unconjugated
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at 4° C prior to opening. This product is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
Synonyms	Dog Immunoglobulin G F(c) fragment, IgG Fc
Application Note	Dog IgG F(c) Fragment can be utilized as a control or standard reagent in Western Blotting and ELISA experiments. Dog IgG F(c) Fragment is stable at 4° C prior to restoration. It is recommended to aliquot restored Dog IgG F(c) Fragment and store at -20° C for extended storage and to prevent repeated freeze-thaw cycles.
Background	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. This product possesses the F(c) region, recognized by high-affinity Fc receptor proteins.
Purity And Specificity	Dog IgG F(c) Fragment was prepared from normal serum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and papain digestion followed by chromatographic separation and extensive dialysis against the buffer stated above. Dog IgG F(c) Fragment was assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Dog Serum, anti-Dog IgG and anti-Dog IgG F(c). No reaction was observed against anti-Dog IgG F(ab') ₂ or anti-Papain.
Assay Dilutions	User Optimized
ELISA	User Optimized
Western Blot	User Optimized
Immunohistochemistry	User Optimized
Other Assays	User Optimized
Expiration	Expiration date is one (1) year from date of opening.

Related Products

010-0102	MOUSE IgG whole molecule - 010-0102
104-4101	Anti-DOG SERUM (RABBIT) Antibody - 104-4101
604-1203	Anti-DOG IgG F(c) (GOAT) Antibody Fluorescein Conjugated - 604-1203
BSA-50	BOVINE SERUM ALBUMIN - Fraction V (Immunoglobulin and Protease Free) - BSA-50

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.