

DOG IgG F(ab')2 fragment (BULK ORDER) - 004-0104

Co	ode: 004-0104	, <u> </u>	X	Size: 2 mg	
Product Description: DOG lgG F(ab')2 fragment (BULK ORDER) - 004-0104					
· · · ·	2.0 mg/mL by UV absorbance at 280 nm				
PhysicalStat	e: Liquid (sterile	: Liquid (sterile filtered)			
Label	Unconjugate	d			
Buffer	0.01 M Sodiu	um Phosphate, 0.15	M Sodium Chloride, pH	17.2	
Stabilizer	None				
Preservative	0.01% (w/v)	Sodium Azide			
Storage Condition	Store vial at 4° C upon receipt. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.				
Synonyms	Dog IgG F(a	b')2 fragment, Dog I	gG Fab2 fragment		
Application Note	Dog IgG F(a	b')2 can be utilized a	as a control or standard	reagent in Western Blotting and ELISA experiments.	
Background	serum immu destruction c cascade, and	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. This product possesses the F(ab')2 fragment, recognized by the two F(ab) fragments yielded from the digestion of the antibody below the disulfide bond hinge region.			
Purity And Specificity	fractionation buffer stated	This product was prepared from normal serum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by pepsin digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Dog IgG, anti-Dog IgG F(ab')2 and anti-Dog Serum. No reaction was observed against anti-Dog IgG F(c) or anti-Pepsin.			
Expiration	Expiration da	Expiration date is one (1) year from date of opening.			
Related Products					
0	10-0102	MOUSE IgG who	ble molecule - 010-0102		
6	10-4302	Anti-MOUSE IgG Conjugated - 610	6 (H&L) (RABBIT) Antibo 0-4302	ody Peroxidase	
6	11-1302	Anti-RABBIT IgG - 611-1302	i (H&L) (GOAT) Antibod	ly Peroxidase Conjugated	
В	SA-50	BOVINE SERUM Protease Free) -	1 ALBUMIN - Fraction V BSA-50	(Immunoglobulin and	
Disalaiman					

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.