

HUMAN IgG whole molecule - 009-0102
Code: 009-0102

Size: 10 mg

Product Description: HUMAN IgG whole molecule - 009-0102

Concentration: 10.0 mg/mL by UV absorbance at 280 nm

PhysicalState: Lyophilized


Label	Unconjugated
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store Human IgG at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Synonyms	Human IgG whole molecule, Human Immunoglobulin G
Application Note	Human IgG whole molecule can be utilized as a control or standard reagent in Western Blotting, Flow, and ELISA experiments.
Background	Human IgG purified protein (Immunoglobulin G) are antibody molecules. Human IgG is composed of four peptide chains — two heavy chains gamma and two light chains. Human IgG has two antigen binding sites. Other Immunoglobulins may be described in terms of polymers with the IgG structure considered the monomer. Human IgG typically constitutes 75% of serum immunoglobulins. Human IgG molecules are synthesized and secreted by plasma B cells.
Purity And Specificity	IgG was prepared from normal human serum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Human IgG and anti-Human Serum.
Assay Dilutions	User Optimized
ELISA	1:10,000 - 1:50,000
Western Blot	1:1,000 - 1:5,000
Flow Cytometry	1:500
Other Assays	User Optimized
Expiration	Expiration date is one (1) year from date of opening.

Related Products

010-0102	MOUSE IgG whole molecule - 010-0102
610-4302	Anti-MOUSE IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated - 610-4302
611-1302	Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302
BSA-50	BOVINE SERUM ALBUMIN - Fraction V (Immunoglobulin and Protease Free) - BSA-50

Images

1	SDS-PAGE of Human IgG. Lane 1: Reduced Human IgG. Lane 2: 5µL OPAL Pre-stained Marker MB-210-0500. Lane 3: Non-reduced Human IgG. Load: 1µg per lane. Predicted/Observed size: Non-reduced at 180-245 kDa, Reduced at 55, 25 kDa.
---	---



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.