

**HUMAN TRANSFERRIN Biotin conjugated - 009-0634**
**Code:** 009-0634

**Size:** 1 mg

**Product Description:** HUMAN TRANSFERRIN Biotin conjugated - 009-0634

**Concentration:** 1.0 mg/mL by UV absorbance at 280 nm

**PhysicalState:** Lyophilized

<b>Label</b>	Biotin
<b>Buffer</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Reconstitution Volume</b>	1.0 mL
<b>Reconstitution Buffer</b>	Restore with deionized water (or equivalent)
<b>Stabilizer</b>	10 mg/ml Polyethylene Glycol (PEG-8000)
<b>Preservative</b>	0.01% (w/v) Sodium Azide
<b>Storage Condition</b>	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
<b>Synonyms</b>	Human Transferrin Biotin Conjugation
<b>Application Note</b>	Human Transferrin Biotin is designed for Immunoblotting, ELISA, immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates.
<b>Background</b>	Human transferrin is encoded by the TF gene and is an iron-binding blood plasma glycoprotein that controls the level of free iron in biological fluids. Human transferrin binds iron very tightly but reversibly. Human transferrin is the most important iron pool in mammals. Human transferrin has a molecular weight of around 80 kDa and contains 2 specific high-affinity Fe(III) binding sites. The affinity of Human transferrin for Fe(III) is extremely high but decreases progressively with decreasing pH below neutrality. Human Transferrin also plays a role in the immune system, creating environments low in iron for which many pathogenic bacteria are unable to thrive.
<b>Purity And Specificity</b>	This product was prepared from normal serum by a multi-stage process that includes delipidation and selective precipitation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Human Transferrin and anti-Human Serum.
<b>Assay Dilutions</b>	User Optimized
<b>Other Assays</b>	User Optimized
<b>Expiration</b>	Expiration date is one (1) year from date of opening.

**Related Products**

010-0102	MOUSE IgG whole molecule - 010-0102
610-4302	Anti-MOUSE IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated - 610-4302
611-1302	Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302
BSA-50	BOVINE SERUM ALBUMIN - Fraction V (Immunoglobulin and Protease Free) - BSA-50

**Related Links**

NCBI - AAB22049.1

<https://www.ncbi.nlm.nih.gov/protein/AAB22049.1>

 UniProtKB - <https://www.uniprot.org/uniprot/Q06AH7>

GeneID - 7018

## Images

1

Dot Blot of Biotin conjugated Human Transferrin. Antigen: Human Transferrin Biotin. Load: 3-fold serial dilution starting at 200 ng. Primary antibody: None. Secondary antibody: Peroxidase streptavidin secondary antibody at 1:40,000 for 30 min at RT. Block: MB-070 for 60 min at RT.

1

200 ng

66.7 ng

22.2 ng

7.4 ng

2.5 ng

2

Dot Blot results of Human Transferrin Biotin Conjugated. Antigen: Human Transferrin Biotin Conjugated. Blot loaded at 3 fold dilution: 1. 100ng, 2. 33.3ng, 3. 11.1ng, 4. 3.70ng, 5. 1.23ng. Blocking: MB-070 Buffer for 30 minutes at RT. Secondary Antibody: Streptavidin-HRP (p/n S000-03) at 1:40,000 for 30 min at RT. Imaging System ChemiDoc, Filter used: Chemi.

1

2

3

4

5

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.