



MOUSE IgG1 isotype control Texas Red™ conjugated - 010-0940

Code: 010-0940

Size: 100 µL

Product Description: MOUSE IgG1 isotype control Texas Red™ conjugated - 010-0940

Concentration: 0.5 mg/mL by UV absorbance at 280 nm

PhysicalState: Lyophilized

| | |
|-------------------------------|---|
| Label | Texas Red® |
| Emission Wavelength | 620 |
| Excitation Wavelength | 596 |
| Buffer | 0.02 M Potassium Phosphate, 0.5 M Sodium Chloride, pH 7.2 |
| Reconstitution Volume | 100 µL |
| Reconstitution Buffer | Restore with deionized water (or equivalent) |
| Stabilizer | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| Preservative | 0.01% (w/v) Sodium Azide |
| Storage Condition | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| Synonyms | Mouse IgG1 subclass isotype control, Mouse IgG1 isotype control |
| Application Note | Mouse IgG1 isotype control can be utilized as a control or standard reagent in Flow cytometry, Western Blotting, and ELISA experiments where determination of sample isotype is important. |
| Background | Isotype control Mouse IgG1 is important for Flow Cytometry. Mouse IgG1 control has no specificity for target cells within a particular experiment. Their purpose is to confirm the specificity of primary antibody binding that it is not a result of non-specific Fc receptor binding to cells or other cellular protein interactions. Isotype controls need to be matched to the specific primary Abs (species and isotype, including heavy and light chains) being used. |
| Purity And Specificity | This product has been prepared from mouse ascitic fluid by immunoaffinity chromatography using protein A. In an Ouchterlony double diffusion assay the material is non-reactive with antisera to mouse IgG2a , IgG2b , IgG3 , IgM , and IgA. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Mouse IgG and anti-Mouse serum. |
| Assay Dilutions | User Optimized |
| FLISA | User Optimized |
| IF Microscopy | 1:1000-1:5000 |
| Flow Cytometry | 1:1000-1:5000 |
| Other Assays | User Optimized |
| Expiration | Expiration date is one (1) year from date of opening. |

Related Products

| | |
|----------|---|
| 010-0102 | MOUSE IgG whole molecule - 010-0102 |
| 610-4302 | Anti-MOUSE IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated - 610-4302 |
| 611-1302 | Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302 |
| BSA-50 | BOVINE SERUM ALBUMIN - Fraction V (Immunoglobulin and Protease Free) - BSA-50 |

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.