

RABBIT IgG F(c) fragment - 011-0103
Code: 011-0103

Size: 1 mg

Product Description: RABBIT IgG F(c) fragment - 011-0103

Concentration: 1.1 mg/mL by UV absorbance at 280 nm

PhysicalState: Liquid (sterile filtered)


Label	Unconjugated
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at 4° C prior to opening. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Rabbit IgG F(c) fragment is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Synonyms	Rabbit immunoglobulin Gamma F(c), IgG Fc
Application Note	Rabbit IgG F(c) Fragment can be utilized as a control or standard reagent in Western Blotting and ELISA experiments. Rabbit IgG F(c) Fragment is stable at 4° C prior to restoration. It is recommended to aliquot restored Rabbit IgG F(c) Fragment and store at -20° C for extended storage and to prevent repeated freeze-thaw cycles.
Background	<p>Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. The F(c) fragment binds with very high affinity to the Fc receptor proteins on phagocytic leukocytes. When digested from the whole antibody molecule, the F(c) fragment no longer possesses the epitope recognition site.</p> <p>Rabbit IgG F(c) fragment is ideal for investigators in Immunology, Cancer, and Microbiology research.</p>
Purity And Specificity	Rabbit IgG F(c) fragment was prepared from normal serum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and papain digestion followed by chromatographic separation and extensive dialysis against the buffer stated above. Rabbit IgG F(c) fragment assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, anti-Rabbit IgG and anti-Rabbit IgG F(c). No reaction was observed against anti-Rabbit IgG F(ab') ₂ or anti-Papain.
Assay Dilutions	User Optimized
ELISA	User Optimized
Western Blot	User Optimized
Immunohistochemistry	User Optimized
Other Assays	User Optimized
Expiration	Expiration date is one (1) year from date of opening.
General Reference	Janeway, Jr., Travers, Walport, and Shlomchik. "The Immune System in Health and Disease." Immunobiology, 5th Edition: Garland Science: 2001.

Related Products

006-0103	GUINEA PIG IgG F(c) fragment - 006-0103(1)
010-0103	MOUSE IgG F(c) fragment - 010-0103
011-0003	RABBIT IgG F(c) fragment Rhodamine conjugated - 011-0003
011-0102	RABBIT IgG whole molecule - 011-0102

Images

SDS-PAGE of Rabbit IgG F(c) Fragment (p/n 011-0103). Lane M: 3 μ L Opal Prestained Marker (p/n MB-210-0500). Lane 1: Reduced Rabbit IgG Whole Molecule (p/n 011-0102). Lane 2: Reduced Rabbit IgG F(ab) Fragment (p/n 011-0105). Lane 3: Reduced Rabbit IgG F(c) Fragment (p/n 011-0103). Lane 4: Reduced Rabbit IgM Whole Molecule (p/n 011-0107). Load: 1 μ g for F(ab) and F(c); 1.2 μ g for IgG and IgM. Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25 kDa; F(ab) at 25 kDa; IgM at 70 and 23 kDa. Observed F(c) Fragment migrates slightly higher.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.