



Mouse Monoclonal Antibody to

EGFR (phospho-Tyr 845)

clone 12A3

biotinylated

Order No.: 0116-100BIOTIN/EGFR-12A3

Size (µg) 100

Lot No.: 0116S



03/230207F

Isotype	Species Reactivity	Applications	Mol. Weight	Ref. Cell Line	Epitope	Immunogen
IgG1	human, mouse	ELISA, WB, IP, IHC (PS, FS), ICC, Luminex	180 kDa	HepG2	Phosphotyrosine 845 E K E pY H A E	phosphopeptide conjugated to KLH

Background and Specificity:

EGFR/erbB receptors are activated upon binding of EGF and EGF-related growth factors such as TGF alpha, beta-cellulin, Hb-EGF, HRG, or NRG. Binding of these ligands leads to receptor homo- and heterodimerization followed by autophosphorylation and activation of downstream signal transduction pathways (MAPK, PI3K/PKB, and STAT). In addition, EGFR becomes fully activated after phosphorylation of Y845 by src family kinases.

Phosphorylation of Y1045 leads to association with cbl and subsequent receptor degradation.

Phosphorylation of S1047 by CamKinase II leads to attenuation of kinase activity; phosphorylation of T654 (by PKC) and T669 (by MAPK, p38) interferes with receptor endocytosis/recycling.

Mab EGFR-12A3 specifically recognizes EGFR phosphorylated at Tyrosine 845 and detects EGFR activation after interaction with src kinases. Mab 12A3 does not crossreact with the highly homologues pTyr 877 of activated erbB2.

Related Products

Blocking peptide for mab EGFR-12A3

#2003-100/EGFR pTyr845

mab to EGFR (C-terminus)

#0007-100/EGFR-13G8

mab to EGFR (cytoplasmic domain)

#0168-100/EGFR-10F4

mab to EGFR (extracellular domain)

#0209-100/EGFR-20E12

mab to EGFR (aa 960 - 980)

#0199-100/EGFR-16F8

mab to EGFR (N-terminus)

#0201-100/EGFR-14C8

mab to phospho-EGFR (pY1045)

#0136-100/EGFR-11C2

mab to phospho-EGFR (pY1068)

#0187-100/EGFR-15A2

mab to phospho-EGFR (pY 1086)

#0188-100/EGFR-8B8

mab to phospho-EGFR (pY 1148)

#0219-100/EGFR-10G12

mab to phospho-EGFR (pY1173)

#0008-100/EGFR-9H2

mab to dephospho-EGFR (Y1173)

#0009-100/EGFR-20G3

mab to phospho-EGFR (pT669)

#0191-100/EGFR-5F10

mab to phospho-EGFR (pT654)

#0138-100/EGFR-3F2

mab to phospho-EGFR (pS1047)

#0107-100/EGFR-1H9

For monoclonal antibodies against erbB2, phospho-erbB2, erbB3 and erbB4, as well as against various EGFR downstream targets, please refer to our website at www.nanotools.de

Purification:	The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography.
Formulation:	liquid; 0.5 mg/ml in PBS/0.09% Na-Azide/PEG and Sucrose
Reconstitution:	
Stability:	Aliquote and freeze in liquid nitrogen; antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.

Avoid repeated freeze / thaw cycles.

Positive Control:	#0813: Cell lysate from EGF-treated HepG2 cells
Immunoblotting:	1 µg/ml for HRPO/ECL detection Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

Immunoprecipitation:	use at 1 - 10 µg per 10 ⁶ vanadate treated A431 cells
Immunocytochemistry	use at 1 - 10 µg/ml
ELISA:	0.1 µg/ml (protein ELISA); capture ELISA: N.D.

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.