

SWINE IgG F(c) fragment Rhodamine conjugated - 014-0003

Code: 014-0003 Size: 1 mg

Product Description: SWINE IgG F(c) fragment Rhodamine conjugated - 014-0003

PhysicalState: Lyophilized

Label Rhodamine (TRITC)

Emission Wavelength 570

Excitation Wavelength 550

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below.

Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.

Swine IgG F(c) fragment Rhodamine conjugation, Swine IgG F(c) fragment TRITC conjugate **Synonyms**

SWINE IgG F(c) fragment Rhodamine conjugate is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex **Application Note**

analysis, including multicolor imaging, utilizing various commercial platforms.

Background Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of

serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. This product possesses the F(c) region, recognized by high-affinity

Fc receptor proteins.

This product was prepared from normal serum by delipidation, salt fractionation, ion exchange chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Swine IgG, anti-Swine IgG F(c) and anti-Purity And Specificity

Swine Serum. No reaction was observed against anti-Swine IgG F(ab')2 or anti-Papain.

Expiration Expiration date is one (1) year from date of opening.

Related Products

010-0102 MOUSE IgG whole molecule - 010-0102

Anti-MOUSE IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated - 610-4302 610-4302

Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated 611-1302

BSA-50 BOVINE SERUM ALBUMIN - Fraction V (Immunoglobulin and

Protease Free) - BSA-50

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.