

## Mouse Monoclonal Antibody to

# MAPK2/erk2 (N-terminus)

## clone 6H3

## biotinylated

**Order No.:** 0178-100BIOTIN/MAPK2-6H3  
**Size (µg)** 100  
**Lot No.:** 0178S



03/150307F

Isotype	Species Reactivity	Applications	Mol. Weight	Ref.Cell Line	Epitope	Immunogen
IgG1	human, mouse, rat, dog	WB, ELISA	42 kDa	HepG2	N-terminus	peptide conjugated to hemocyanin

### Background and Specificity:

Extracellular signal/mitogen activated protein kinases (erk/MAPK) are a group of proline-directed serine/threonine kinases that are activated by dual phosphorylation of conserved threonine and tyrosine residues within a characteristic **T X Y** peptide motif. The mitogen-activated kinases erk1 (MAPK1) and erk2 (MAPK2) acquire full enzymatic activity upon phosphorylation of both threonine and tyrosine residues within the sequence motif **T E Y**.

**Mab MAPK2-6H3** specifically recognizes the N-terminus of MAP kinase 2 (erk2).

### Related Products

- mab to MEK1 (pS218/222)**
- mab to MEK2 (pS222/226)**
- #0174-100/MEK1/2-7E10
- mab to MEK1 (N-terminus)**
- #0186-100/MEK1.10B1
- mab to MEK1/2**
- #0150-100/MEK1/2-9G3
- mab to MEK2 (N-terminus)**
- #0148-100/MEK2-8E8
- mab to MKK3 (N-terminus)**
- #0166-100/MKK3-5F7
- mab to MKK5 (N-terminus)**
- #0224-100/MKK5-14B5
- mab to MKK7 (N-terminus)**
- #0189-100/MKK7-10F7
- mab to MAPK 1/2 (pT-E-pY)**
- #0012-100/MAPK-12D4
- mab to MAPK 2 (C-terminus)**
- #0011-100/MAPK-6G11
- mab to MAPK 2 (internal sequence)**
- #0239-100/MAPK2-12A4
- mab to MAPK7/erk5**
- #0223-100/MAPK7/erk5-12F2
- mab to Fos (pS374)**
- #0118-100/Fos-34E4
- mab to Fos (N-terminus)**
- #0122-100/Fos-8B5
- mab to C-Raf (pS621)**
- #0102-100/C-Raf-6B4
- mab to C-Raf**
- #0120-100/C-Raf-PBB-1

**Purification:** The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography.

**Formulation:** liquid, 0.5 mg/ml in PBS/0,09% Na-Azide/PEG and Sucrose

**Reconstitution:**

**Stability:** Aliquote and freeze in liquid nitrogen; reconstituted antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.

**Avoid repeated freeze / thaw cycles.**

**Positive Control:** #0811: Cell lysate from untreated HepG2 cells

**Immunoblotting:** 0.5 µg/ml for HRPO/ECL detection  
**Recommended blocking buffer:** Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

**Immunoprecipitation:** ND

**Immunocytochemistry:** ND

**ELISA:** use at 0.05 µg/ml

**All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.**