

Mouse Monoclonal Antibody to

EGFR (phospho-Tyr 1086)

clone 8B8

biotinylated

Order No.: 0188-100BIOTIN/EGFR-8B8

Size (µg) 100

Lot No.: 0188S

02/260207F

Isotype	Species Reactivity	Applications	Mol. Weight	Ref. Cell Line	Epitope	Immunogen
IgG2b	human	WB, ELISA	180 kDa	A431	phosphotyrosine 1086 N P V p Y H N Q	synthetic phosphopeptide conjugated to hemocyanin

Background and Specificity:

EGFR/erbB receptors are activated upon binding of EGF and EGF-related growth factors such as TGF alpha, beta-cellulin, Hb-EGF, HRG, or NRG. Binding of these ligands leads to receptor homo- and heterodimerization followed by autophosphorylation and activation of downstream signal transduction pathways (MAPK, PI3K/PKB, and STAT). In addition, EGFR becomes fully activated after phosphorylation of Y845 by src family kinases.

Phosphorylation of Y1045 leads to association with cbl and subsequent receptor degradation.

Phosphorylation of S1047 by CamKinase II leads to attenuation of kinase activity; phosphorylation of T654 (by PKC) and T669 (by MAPK, p38) interferes with receptor endocytosis/recycling.

Mab EGFR-8B8 specifically recognizes EGFR phosphorylated at Tyrosine 1086. The antibody does not interact with the non-phosphorylated EGFR nor with unrelated Tyrosine-phosphorylated proteins.

Related Products

mab to EGFR (C-terminus)
#0007-100/EGFR-13G8
mab to EGFR (cytoplasmic domain)
#0168-100/EGFR-10F4
mab to EGFR (extracellular domain)
#0209-100/EGFR-20E12
mab to EGFR (aa 960 - 980)
#0199-100/EGFR-16F8
mab to EGFR (N-terminus)
#0201-100/EGFR-14C8
mab to phospho-EGFR (pY 845)
#0116-100/EGFR-12A3
mab to phospho-EGFR (pY1045)
#0136-100/EGFR-11C2
mab to phospho-EGFR (pY1068)
#0187-100/EGFR-15A2
mab to phospho-EGFR (pY 1148)
#0219-100/EGFR-10G12
mab to phospho-EGFR (pY1173)
#0008-100/EGFR-9H2
mab to dephospho-EGFR (Y1173)
#0009-100/EGFR-20G3
mab to phospho-EGFR (pT669)
#0191-100/EGFR-5F10
mab to phospho-EGFR (pT654)
#0138-100/EGFR-3F2
mab to phospho-EGFR (pS1047)
#0107-100/EGFR-1H9

For monoclonal antibodies against erbB2, phospho-erbB2, erbB3 and erbB4, as well as against various EGFR downstream targets, please refer to our website at www.nanotools.de

Purification: The antibody was purified from serum-free cell culture supernatant by subsequent ultrafiltration and size exclusion chromatography.

Formulation: liquid; 0.5 mg/ml in PBS/0.09% Na-Azide/PEG and Sucrose

Reconstitution:

Stability: Aliquote and freeze in liquid nitrogen; antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.

Avoid repeated freeze / thaw cycles.

Positive Control: #0832: Cell lysate from pervanadate-treated A431 cells

Immunoblotting: 1 µg/ml for HRPO/ECL detection
Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

Immunoprecipitation: ND

Immunocytochemistry: ND

ELISA: use at 0.1 µg/ml

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.