Fax: Mouse Monoclonal Antibody to

SAPK1α/jnk2 (N-Terminus)





biotinylated clone 12C5 0190-100BIOTIN/SAPK1a/jnk2-12C5 Order No.: Size (µg) 100 0190S Lot No.: 04/080507F Mol. Weight Isotype **Species Reactivity** Applications **Ref.Cell Line** Epitope Immunogen human, mouse, rat, WB, ELISA, 54 kDa A431 N-terminus peptide conjugated to IHC hemocyanin dog **Related Products Background and Specificity:** mab to SAPK1β/jnk3 (N-terminus) Stress-activated Protein Kinases (SAPKs) are strongly activated in response to adverse stimuli such as heat and osmotic shock, UV light and other DNA-damaging reagents, and inhibitors of #0220-100/SAPK1b/jnk3-4G6 mab to SAPK1y/jnk1 (N-terminus) protein synthesis. They are also activated strongly in response to agonists that are released or #0200-100/SAPK1a/ink1-5D10 produced under conditions of stress, such as proinflammatory cytokines. mab to SAPK1/2 (T - G/P - pY) #0041-100/SAPK1/2-9H The SAPK1/jnk family consists of 3 isoforms: SAPK1 α /jnk2; SAPK1 β /jnk3; SAPK1 γ /jnk1. mab to SAPK2a (N-terminus) #0034-100/SAPK2a-13D5 (crossreaction with Mxi2) #0035-100/SAPK2a-20B11 (no crossreaction with Mxi2) Mab SAPK1a-12C5 specifically interacts with the N-terminus of SAPK1a/jnk2 kinases. mab to SAPK28 (N-terminus) #0053-100/SAPK2d-5H7 mab to MAPK 1/2 (pT-E-pY) #0012-100/MAPK-12D4 mab to MAPK 2 (C-terminus) #0011-100/MAPK2-6G mab to MAPK 2 (N-terminus) #0178-100/MAPK2-6 mab to MAPK 2 (internal sequence) #0239-100/MAPK2-12 **Purification:** The antibody was purified from serum-free cell culture mab to MAPK7/erk5 supernatant by subsequent thiophilic adsorption and size #0223-100/MAPK7/erk5-12F2 exclusion chromatography. mab to Mxi 2 (N-terminus) 0046-100/Mxi-2F2 liquid; 0.5 mg/ml in PBS/0.09% Na-Azide/PEG and Sucrose Formulation: **Reconstitution:** Stability: Aliquote and freeze in liquid nitrogen; antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months. Avoid repeated freeze / thaw cycles. **Positive Control:** #0831: Cell lysate from untreated A431 cells Immunoblotting: 0.5 µg/ml for HRPO/ECL detection Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT. Immunoprecipitation: ND ND Immunocytochemistry: use at 0.05 µg/ml All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.

lgG1

ELISA: