

Mouse Monoclonal Antibody to

LC3 (microtubule-associated protein1 light chain 3B) clone 5F10 biotinylated



Order No.: 0231-100BIOTIN/LC3-5F10
Size (µg) 100
Lot No.: 0231S

06/270208F

Isotype	Species Reactivity	Applications	Mol. Weight	Ref.Cell Line	Epitope	Immunogen
IgG1	human, mouse, rat, dog, hamster	ICC, WB	LC3-I: 18kDa LC3-II:16kDa	Neuro 2A	N-terminus of LC3-B	synthetic peptide conjugated to hemocyanin

Background and Specificity:

Autophagy is an alternative process of proteasomal degradation for some long-lived proteins or organelles. Alterations in the autophagic-lysosomal compartment have been linked to neuronal death in many neurodegenerative disorders as well as in transmissible neuronal pathologies (prion diseases). Genetic studies in yeast have shown that Autophagy-defective Gene-8 (Atg-8) represents a specific marker for autophagy. Among the four families of mammalian Atg8-related proteins only LC3 (microtubule-associated protein1 light chain 3) is expressed at sufficient high levels and efficiently recruited to autophagic vesicles in cells and tissues. During autophagy the cytoplasmic form, LC3-I is processed and recruited to autophagosomes, where LC3-II is generated by site specific proteolysis near to the C-terminus. Autophagic vacuoles have been also reported frequently in cardiomyopathies or muscle cells exposed to different experimental settings.

Mab LC3-5F10 specifically recognizes both forms of endogenous LC3, the cytoplasmic LC3-I (18 kDa) as well as the lipidated form generated during autophagosome and autophagolysosome formation: LC3-II (16 kDa). Immunocytochemical staining of cells with LC3-5F10 mab reveals the specific punctate distribution of endogenous LC3-II as a hallmark of autophagic activity.

Related Products

- mab to LC3**
#0260-100/LC3-2G6
- mab to LC3**
#0261-100/LC3-5H3
- mab to Beclin**
#0240-100/Beclin-12B4
- Alzheimer Disease**
- mab to βA4 (1-40), C-Terminus**
#0060-100/bA4(40)-5C3
- mab to βA4 (1-42), C-Terminus**
#0061-100/bA4(42)-8G7
- mab to βA4 (1-40/42), C-Terminus**
#0062-100/bA4(40/42)-9F1
- mab to βA4 (1-43), C-Terminus**
#0095-100/bA4(43)-6G12
- mab to βA4, N-Terminus**
#0064-100/bA4N-19H5
- mab to βA4, N-Terminus**
#0084-100/bA4N-19H11
- mab to βA4, N-Terminus**
#0197-100/bA4N-11H3

For monoclonal antibodies against PKB/akt, and SAPK/jnk, please refer to our website at www.nanotools.de

Purification:	The antibody was purified from serum-free cell culture supernatant by subsequent ultrafiltration and size exclusion chromatography.
Formulation:	liquid in PBS / 0.09 % Na-azide / PEG and Sucrose.
Reconstitution:	
Stability:	Aliquote and freeze in liquid nitrogen; antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months. Avoid repeated freeze / thaw cycles
Positive Control:	#0911: Cell lysate from untreated Neuro 2A
Immunoblotting:	0.5 µg/ml for HRPO/ECL detection Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.
Immunoprecipitation:	ND
Immunocytochemistry:	Use at 1- 10 µg/ml (paraformaldehyd/methanol fixation)
ELISA:	ND

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.