

## Anti-PEROXIDASE (Horseradish) (MOUSE) Antibody - 100-3138

**Code:** 100-3138

**Size:** 2 mL

**Product Description:** Anti-PEROXIDASE (Horseradish) (MOUSE) Antibody - 100-3138

**Concentration:** 85 mg/mL by Refractometry

**PhysicalState:** Lyophilized

<b>Label</b>	Unconjugated
<b>Host</b>	Mouse
<b>Buffer</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Reconstitution Volume</b>	2.0 mL
<b>Reconstitution Buffer</b>	Restore with deionized water (or equivalent)
<b>Stabilizer</b>	None
<b>Preservative</b>	0.01% (w/v) Sodium Azide
<b>Storage Condition</b>	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
<b>Synonyms</b>	Mouse Anti-PEROXIDASE Antibody, Anti-PEROXIDASE Horseradish, mouse anti-HRP antibody, mouse anti Peroxidase, mouse anti HRP
<b>Application Note</b>	Suitable for immunoblotting (western or dot blot), ELISA, immunoprecipitation and most immunological methods requiring high titer and specificity.
<b>Background</b>	Peroxidase antibody detects lipid peroxides. Peroxidases are a large family of enzymes that typically catalyze a reaction of the form: ROOR' + electron donor (2 e-) + 2H+ ROH + R'OH. For many of these enzymes the optimal substrate is hydrogen peroxide, but others are more active with organic hydroperoxides such as lipid peroxides. Peroxidases can contain a heme cofactor in their active sites, or alternately redox-active cysteine or selenocysteine residues. The nature of the electron donor is very dependent on the structure of the enzyme. For example, horseradish peroxidase can use a variety of organic compounds as electron donors and acceptors. Horseradish peroxidase has an accessible active site, and many compounds can reach the site of the reaction. Because there is a very closed active site, for an enzyme such as cytochrome c peroxidase, the compounds that donate electrons are very specific. While the exact mechanisms have yet to be elucidated, peroxidases are known to play a part in increasing a plant's defenses against pathogens.[1] Peroxidases are sometimes used as histological marker. Cytochrome c peroxidase is used as a soluble, easily purified model for cytochrome c oxidase. Anti-Peroxidase antibody is ideal for investigators involved in enzyme research.
<b>Purity And Specificity</b>	Anti-Peroxidase antibody was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-mouse serum, purified and partially purified Peroxidase [Horseradish]. Cross reactivity against Peroxidase from other tissues and species may occur but have not been specifically determined.
<b>ELISA</b>	1:5,000 - 1:20,000
<b>Western Blot</b>	1:500 - 1:2,000
<b>Immunohistochemistry</b>	User Optimized
<b>Expiration</b>	Expiration date is one (1) year from date of opening.
<b>Immunogen</b>	Peroxidase [Horseradish]
<b>General Reference</b>	Karthikeyan M et al. (December 2005). "Induction of resistance in host against the infection of leaf blight pathogen ( <i>Alternaria palandui</i> ) in onion ( <i>Allium cepa</i> var <i>aggregatum</i> )". <i>Indian J Biochem Biophys</i> 42 (6): 371-7.

### Related Products

200-301-268	Anti-AKT pS473 (MOUSE) Monoclonal Antibody - 200-301-268
610-4302	Anti-MOUSE IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated - 610-4302
611-1302	Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302

FEMTOMAX-020 Chemiluminescent FemtoMax™ Super Sensitive HRP Substrate  
for Microwell and/or Membrane (2 component system) -  
FEMTOMAX-020

#### **Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.