

Anti-IKK β (RABBIT) Antibody - 100-401-220

Code: 100-401-220

Size: 100 μ L

Product Description: Anti-IKK β (RABBIT) Antibody - 100-401-220

Concentration: 80 mg/mL by Refractometry

PhysicalState: Liquid (sterile filtered)

Label	Unconjugated
Host	Rabbit
Gene Name	IKBKB, IKKB
Species Reactivity	human, mouse, rat
Buffer	None
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Synonyms	rabbit anti-IKK beta antibody, Inhibitor of nuclear factor kappa-B kinase subunit beta, I-kappa-B-kinase beta, IkbKB, IKK-beta, IKK-B, I-kappa-B kinase 2, IKK2, Nuclear factor NF-kappa-B inhibitor kinase beta, NFKBIKB
Application Note	This product was assayed by immunoblot and found to be reactive against IKK β at a dilution of 1:1000 followed by reaction with Peroxidase conjugated Affinity Purified anti-Rabbit IgG [H&L] (Goat) code #611-1302. Anti-IKK β is suitable for the detection by immunoblot of human, mouse and rat IKK β showing an 87 kDa band.
Background	NF κ B comprises a family of cellular transcription factors that are involved in the inducible expression of a variety of cellular genes that regulate the inflammatory response and control of cell death. In the cytoplasm NF κ B is negatively modulated by the inhibitory proteins I κ B. In turn I κ B is phosphorylated by a cellular kinase complex called IKK. IKK is a heterodimer composed of two kinases: IKK-a and IKK-b that phosphorylate I κ B leading to its degradation and the resulting translocation of NF κ B to the nucleus. IKK kinase activity is modulated negatively by pharmaceutical agents such as aspirin and positively by various cellular components such as TNF- α , endotoxins and overexpression of cellular kinases like MEK1. Aspirin appears to have its effect by inhibiting the binding of ATP to IKK.
Purity And Specificity	This product was prepared from monospecific antiserum by delipidation and defibrination. Anti- IKK β may react non-specifically with other proteins. Control peptide (code #100-401-220p) will compete only with the specific reaction of antiserum with the IKK β subunit.
Assay Dilutions	User Optimized
ELISA	1:1,000 - 1:5,000
Western Blot	1:200 - 1:1,000
Immunohistochemistry	1:200 - 1:1,000
Other Assays	User Optimized
Expiration	Expiration date is one (1) year from date of opening.
Immunogen	IKK β peptide corresponding to the highly conserved C-terminus region of the human protein conjugated to Keyhole Limpet Hemocyanin (KLH).

Related Products

100-401-219	Anti-IKK α (RABBIT) Antibody - 100-401-219
100-401-401	Anti-AKT (RABBIT) Antibody - 100-401-401
100-4167C	Anti-IKBa C-terminal (RABBIT) Antibody - 100-4167C
100-4186	Anti-IKKBß (RABBIT) Antibody - 100-4186

Related Links

UniProtKB - O14920

<http://www.uniprot.org/uniprot/O14920>


NCBI - O14920.1 <http://www.ncbi.nlm.nih.gov/protein/O14920.1>

GenID - 3551

Images


1

Western blot of anti-IKK β antibody reacted with HeLa cell extract. All incubations except color development were performed using TBS supplemented with 0.1% Tween-20 at room temperature. The membrane was blocked in 5% dry milk for 2 h. After washing, a 1:500 dilution of the primary antibody was added to the membrane and incubated for 2 h. Washes with buffer were performed 4 times for 5' each. The western blot was incubated with secondary antibody (HRP Goat-a-Rabbit IgG [H&L]) diluted 1:2,000 for 1h. Washes with TBS preceded color development.



2

Rockland's Anti-IKK β antibody was diluted 1:500 to detect IKK β in human placenta tissue. Tissue was formalin fixed and paraffin embedded. No pre-treatment of sample was required. The image shows the localization of antibody as the precipitated red signal, with a hematoxylin purple nuclear counter stain.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.