



Anti-ESTERASE (RABBIT) Antibody - 100-4169

Code: 100-4169

Size: 2 mL

Product Description: Anti-ESTERASE (RABBIT) Antibody - 100-4169

Concentration: 85 mg/mL by Refractometry

PhysicalState: Lyophilized

Label	Unconjugated
Host	Rabbit
Gene Name	CES3
Species Reactivity	swine
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Volume	2.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Synonyms	rabbit anti-Esterase Antibody, Liver carboxylesterase, Proline-beta-naphthylamidase, Retinyl ester hydrolase
Application Note	This product has been assayed against 1.0 ug of Esterase [Porcine Liver] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Rabbit IgG [H&L] (Goat) code #611-1302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:8,000 to 1:40,000 of the reconstitution concentration is suggested for this product.
Background	Esterase is involved in the detoxification of xenobiotics and in the activation of ester and amide prodrugs. It is active towards triacylglycerides containing short-chain fatty acids from C2 to C6, and 13-monoacylglycerols containing fatty acids from C2 to C12; and inactive on long-chain triacylglycerols and diacylglycerol. It hydrolyzes aromatic and alkyl esters and vitamin A acetate. The hydrolysis rate depends upon the amino acid promoiety and the esterification site of the prodrug. Aromatic promoieties are favored, highest rates are observed with phenylalanyl prodrugs, hydrolysis of valyl and isoleucyl prodrugs is less efficient. With floxuridine prodrugs, activity is higher on 5' monoesters than on 3' monoesters; while with gemcitabine prodrugs, activity is higher on 3' monoesters than on 5' monoesters.
Purity And Specificity	This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-rabbit serum, purified and partially purified Esterase [Porcine Liver]. Cross reactivity against Esterase from other tissues and species may occur but have not been specifically determined.
Assay Dilutions	User Optimized
ELISA	1:5,000 - 1:25,000
Western Blot	1:500 - 1:3,000
Other Assays	User Optimized
Expiration	Expiration date is one (1) year from date of opening.
Immunogen	Esterase [Porcine Liver]
Related Products	
611-1302	Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302
B304	NORMAL GOAT SERUM (NGS) - B304

FEMTOMAX-110 Chemiluminescent FemtoMax™ Super Sensitive HRP Substrate for Microwell and/or Membrane (2 component system) - FEMTOMAX-110

MB-070 Blocking Buffer for Fluorescent Western Blotting - MB-070

Related Links

UniProtKB - Q29550

<http://www.uniprot.org/uniprot/Q29550>

NCBI - 343791015 <http://www.ncbi.nlm.nih.gov/protein/343791015>

GeneID - 100626873

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.