



Anti-Swine IL-13 (RABBIT) Antibody - 200-401-B55

Code: 200-401-B55

Size: 100 µg

Product Description: Anti-Swine IL-13 (RABBIT) Antibody - 200-401-B55

Concentration: 1.0 mg/mL by UV absorbance at 280 nm

PhysicalState: Lyophilized

Label	Unconjugated
Host	Rabbit
Gene Name	IL-13
Species Reactivity	swine, sheep, bovine
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Volume	100 µL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Synonyms	rabbit anti-Interleukin-13 antibody, rabbit anti-IL-13 antibody, IL13, Interleukin13, IL 13, Interleukin 13
Application Note	IL-13 antibody has been tested for use in ELISA and western blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 13.2 kDa in size corresponding to swine IL-13 protein by western blotting in the appropriate cell lysate or extract.
Background	Interleukin 13 (IL-13), of the IL-4 superfamily, is a cytokine secreted by many cell types, but especially T helper type 2 (Th2) cells, that is an important mediator of allergic inflammation and disease. IL-13 induces its effects through a multi-subunit receptor that includes the alpha chain of the IL-4 receptor (IL-4R) and at least one IL-13-specific binding chain. Most of the biological effects of IL-13, like those of IL-4, are linked to a single transcription factor, STAT6. In humans, IL-13 can induce immunoglobulin E (IgE) secretion from activated B cells. In mice, deletion of IL-13 does not markedly affect either Th2 cell development or antigen-specific IgE responses induced by potent allergens. Deletion of IL-4 abrogates these responses. IL-13 acts as a molecular bridge linking allergic inflammatory cells to the non-immune cells in contact with them, thus altering physiological function. Although IL-13 is associated primarily with the induction of airway disease, including airway hyperresponsiveness, goblet cell metaplasia and mucus hypersecretion, it also induces airway matrix metalloproteinases as part of a mechanism that protects against excessive allergic inflammation that predisposes to asphyxiation. Anti-IL-13 antibody is ideal for investigators involved in Cancer and Immunology research.
Purity And Specificity	This product was Protein-A purified from monospecific antiserum by chromatography. This antibody is specific for swine IL-13 protein. A BLAST analysis was used to suggest reactivity with IL-13 from swine sources based on 100% homology with the immunizing sequence. The following homologies may indicate chances of cross-reactivity when amino acid similarity and not just exact sequence is accounted: 89-90% homology to IL-13 from sheep, dolphin, beluga, and bovine, 83-88% to human and certain monkey IL-13, 80-82% to dusky titi, macaque, baboon, mangabey, dog, llama, camel, and galago, 75-78% to horse, and 67-72% to rat, mouse, and gerbil. Cross-reactivity with IL-13 from other sources has not been determined.
Assay Dilutions	User Optimized
ELISA	1:11,000
Western Blot	1.5 to 2 µg/ml
Other Assays	User Optimized
Expiration	Expiration date is one (1) year from date of opening.
Immunogen	This protein-A purified antibody was prepared from whole rabbit serum produced by repeated immunizations with full length recombinant protein raised in yeast, corresponding to mature swine IL-13 protein.

General Reference

Wendy L. Trigona, Wendy C. Brown, and D. Mark Estes (1999). Functional implications for signaling via the IL4R/IL13R complex on bovine cells. *Veterinary Immunology and Immunopathology* 72(1-2), 73-79
doi:10.1016/S0165-2427(99)00117-8. Lew, D.Lyles, K.Dempsey, Y.Zhao (2000) Mitogenic effect of IL-13 in bovine airway smooth muscle cells (ASMC). *Journal of Allergy and Clinical Immunology*, Volume 105, Issue 1, Page S299

Related Products

200-401-B55S	Anti-Swine IL-13 (RABBIT) Antibody - 200-401-B55S
200-401-B61	Anti-Swine CCL3L1 (RABBIT) Antibody - 200-401-B61
214-401-306	Anti-Swine TNFa (RABBIT) Antibody - 214-401-306
214-401-B85	Anti-Swine MIP-1ß (RABBIT) Antibody - 214-401-B85

Related Links

UniProtKB - B3GDZ5


<http://www.uniprot.org/uniprot/B3GDZ5>

NCBI - 189176151 <http://www.ncbi.nlm.nih.gov/protein/189176151>

GeneID - 396721

Images

- 1 Western blot using Rockland's protein-A purified anti-swine IL-13 antibody shows detection of recombinant swine IL-13 at 13.2kDa (arrow) raised in yeast. Multiple bands are expected of the glycosylated protein. Protein was purified and resolved by SDS-PAGE, then transferred to PVDF membrane. Membrane was blocked with 3% BSA (BSA-30, diluted 1:10), and probed with Rockland's, Inc. Anti-swine IL-13. After washing, membrane was probed with Dylight™ 649 Conjugated Anti-Rabbit IgG (H&L) (Donkey) Antibody (611-743-127).



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.