



Anti-CATALASE (Bovine Liver) (RABBIT) Antibody Biotin Conjugated - 200-4651S

Code: 200-4651S

Size: 25 µL

Product Description: Anti-CATALASE (Bovine Liver) (RABBIT) Antibody Biotin Conjugated - 200-4651S

Concentration: 1.0 mg/mL by UV absorbance at 280 nm

PhysicalState: Liquid (sterile filtered)

Label	Biotin
Host	Rabbit
Gene Name	CAT
Species Reactivity	bovine
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Synonyms	rabbit anti-Catalase Antibody biotin Conjugation, biotin Conjugated rabbit anti-Catalase Antibody, Cas 1 antibody, CAT antibody, Cs1 antibody, MGC138422 antibody, MGC138424 antibody
Application Note	Anti-Catalase has been assayed against 1.0 ug of Catalase in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenzothiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.
Background	Anti-Catalase antibody serves to protect cells of aerobically respiring organisms from the negative effects of hydrogen peroxide. Additionally Catalase promotes cell growth including T-cells, B-cells, myeloid leukemia, melanoma, mastocytoma, and normal and transformed fibroblast cells. Belonging to the catalase family, Anti-Catalase antibody is ideal for researchers invested in Metabolism, Cancer , Cardiovascular , Signal Transduction, and Tags & Cell Marker research.
Purity And Specificity	Anti-Catalase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Rabbit Serum as well as purified and partially purified Catalase [Bovine Liver]. Cross reactivity against Catalase from other sources may occur but has not been specifically determined.
Assay Dilutions	User Optimized
ELISA	1:10,000 - 1:40,000
Western Blot	1:500- 1:2,500
Other Assays	User Optimized
Expiration	Expiration date is one (1) year from date of opening.
Immunogen	Catalase [Bovine Liver]
Related Products	
610-4302	Anti-MOUSE IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated - 610-4302
611-1302	Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302
B501-0500	BLOTTO Immunoanalytical Grade (Non-Fat Dry Milk) - B501-0500
KCB002	HRP Western Blot Anti-Mouse IgG Antibody - KCB002

Related Links

UniProtKB - P00432


<http://www.uniprot.org/uniprot/P00432>

NCBI - AAI03067.1 <http://www.ncbi.nlm.nih.gov/protein/AAI03067.1>

GeneID - 531682

Images

1 Western Blot of Rabbit anti-Catalase antibody. Lane R: Reduced samples of purified Catalase. Lane NR: Non-reduced samples of purified Catalase. Load: ~1ug of protein per lane. Primary antibody: Catalase antibody at 1:1000 for overnight at 4°C. Secondary antibody: Dylight488™ rabbit secondary antibody at 1:10,000 for 1.5 hrs at RT. Block: MB-070 overnight at 4°C. Predicted/Observed size: 59.9 kDa, 55 kDa for catalase. Other band(s): none.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.