

Anti-DONKEY IgG (H&L) (RABBIT) Antibody Fluorescein Conjugated - 216-4202

Code: 216-4202

Size: 20 mg

Product Description: Anti-DONKEY IgG (H&L) (RABBIT) Antibody Fluorescein Conjugated - 216-4202

Concentration: 10.0 mg/mL by UV absorbance at 280 nm

PhysicalState: Lyophilized

| | |
|-------------------------------|---|
| Label | Fluorescein (FITC) |
| Host | Rabbit |
| Emission Wavelength | 528 |
| Excitation Wavelength | 495 |
| Species Reactivity | Donkey |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reconstitution Volume | 2.0 mL |
| Reconstitution Buffer | Restore with deionized water (or equivalent) |
| Stabilizer | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| Preservative | 0.01% (w/v) Sodium Azide |
| Storage Condition | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| Synonyms | Rabbit anti-Donkey IgG FITC conjugated Antibody, Rabbit anti-Donkey IgG Fluorescein Conjugated antibody |
| Application Note | This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. |
| Background | Anti-Donkey IgG Fluorescein Antibody generated in rabbit detects donkey IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the complement cascade, and opsinization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present. |
| Purity And Specificity | This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-fluorescein, anti-Rabbit Serum, Donkey IgG and Donkey Serum. |
| Assay Dilutions | FLOW CYTOMETRY 1:500 - 1:2,500 |
| FLISA | 1:10,000 - 1:50,000 |
| IF Microscopy | 1:1,000 - 1:5,000 |
| Flow Cytometry | 1:500 - 1:2,500 |
| Other Assays | FLOW CYTOMETRY 1:500 - 1:2,500 |
| Expiration | Expiration date is one (1) year from date of opening. |
| Immunogen | Donkey IgG whole molecule |
| Related Products | |

610-4302 Anti-MOUSE IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated - 610-4302

| | |
|----------|--|
| 611-1302 | Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302 |
| B304 | NORMAL GOAT SERUM (NGS) - B304 |
| BSA-50 | BOVINE SERUM ALBUMIN - Fraction V (Immunoglobulin and Protease Free) - BSA-50 |

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.