

Fibroblast Growth Factor basic, human recombinant (rHuFGF-basic)

Catalog No: 50361
Lot No: XXXXX
Source: *E. coli*
Synonyms: Prostatropin, HBGH-2, HBGF-2, FGF-2, FGF-b

Background

Basic fibroblast growth factor is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein functions as a modifier of endothelial cell migration and proliferation, as well as an angiogenic factor. It acts as a mitogen for a variety of mesoderm- and neuroectoderm-derived cells in vitro, thus is thought to be involved in organogenesis. Three alternatively spliced variants encoding different isoforms have been described. The heparin-binding growth factors are angiogenic agents in vivo and are potent mitogens for a variety of cell types in vitro. There are differences in the tissue distribution and concentration of these 2 growth factors.

Description

Fibroblast Growth Factor-2 (FGF-2) human recombinant produced in *E. coli* is a single, non-glycosylated, polypeptide chain containing 154 amino acids and having a molecular mass of 17.2 kDa. FGF-b is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

The protein was lyophilized from a 0.2 µm filtered solution in 20 mM Tris-HCl, pH 7.4 and 1M NaCl.

Solubility

It is recommended to reconstitute the lyophilized FGF-b in sterile 18 MΩ-cm H₂O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized Fibroblast Growth Factor-2, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution FGF-b should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity


Greater than 98.0% as determined by SDS-PAGE.

Amino Acid Sequence

AAGSITTLPA LPEDGGSGAF PPGHFKDPKR LYCKNGGFLL RIHPDGRVDG VREKSDPHIK LQLQAEERGV VSIKGVCANR
YLAMKEDGRL LASKCVTDEC FFFERLESNN YNTYRSRKYT SWYVALKRTG QYKLGSKTGP GQKAILFLPM SAKS

Activity

The ED₅₀, calculated by the dose-dependant proliferation of murine balb/c 3T3 cells is <0.1 ng/ml, corresponding to a specific activity of 1.0 x 10⁷ units/mg.



Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.