

## Tumor Necrosis Factor alpha, human recombinant (rHuTNF-a)

**Catalog No:** 50435  
**Lot No:** XXXXX  
**Source:** *E. coli*  
**Synonyms:** TNF-alpha, Tumor necrosis factor ligand superfamily member 2, TNF-a, Cachectin, DIF, TNFA, TNFSF2

### Background

Tumor necrosis factor is a cytokine involved in systemic inflammation and is a member of a group of cytokines that all stimulate the acute phase reaction. TNF is mainly secreted by macrophages. TNF causes apoptotic cell death, cellular proliferation, differentiation, inflammation, tumorigenesis and viral replication, TNF is also involved in lipid metabolism, and coagulation. TNF's primary role is in the regulation of immune cells. Dysregulation and, in particular, overproduction of TNF have been implicated in a variety of human diseases- autoimmune diseases, insulin resistance, and cancer.

### Description

Tumor Necrosis Factor-a human recombinant produced in *E. coli* is a single, non-glycosylated, polypeptide chain containing 158 amino acids (157 amino acids of the mature human TNF-alpha and an N-terminal methionine) and having a molecular mass of 17.5 kDa. TNF-alpha is purified by standard chromatographic techniques.

### Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

### Formulation

TNF-alpha Human was lyophilized from a concentrated 1 mg/ml solution containing 20 mM PB, pH-7.2, and 100 mM NaCl.

### Solubility

It is recommended to reconstitute the lyophilized TNF-a in sterile 18 MΩ-cm H<sub>2</sub>O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

### Stability

Lyophilized Tumor Necrosis Factor-a, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution TNF-a should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

### Purity


Greater than 95.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

### Amino Acid Sequence

MVRSSSRTPS DKPVAHVVAN PQAEGQLQWL NRRANALLAN GVELRDNQLV VPSEGLYLIY SQVLFGKGQC PSTHVLTLHT  
 ISRIAVSYQT KVNLLSAIKS PCQRETPEGA EAKPWYEPIY LGGVFQLEKG DRLSAEINRP DYLDFAESGQ VYFGIIAL

### Activity

The specific activity is  $\geq 5.0 \times 10^7$  IU/mg as determined by the cytotoxicity of murine L929 cells in the presence of Actinomycin D.



### Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.