

Fibroblast Growth Factor acidic, human recombinant (rHuFGF-acidic)

Catalog No: 50440
Lot No: XXXXX
Source: *E. coli*
Synonyms: HBGF-1, ECGF-beta, FIBP, FGFIBP, FIBP-1, ECGF, ECGFA, GLI0703, FGF1, FGF-a

Background

Acidic fibroblast growth factor is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein functions as a modifier of endothelial cell migration and proliferation, as well as an angiogenic factor. It acts as a mitogen for a variety of mesoderm- and neuroectoderm-derived cells in vitro, thus is thought to be involved in organogenesis. Three alternatively spliced variants encoding different isoforms have been described. The heparin-binding growth factors are angiogenic agents in vivo and are potent mitogens for a variety of cell types in vitro. There are differences in the tissue distribution and concentration of these 2 growth factors.

Description

Fibroblast Growth Factor-acidic (FGF-1) human recombinant produced in *E. coli* is a single, non-glycosylated, polypeptide chain containing 140 amino acids and having a molecular mass of approximately 15.8 kDa. FGF acidic is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

The protein was lyophilized from a concentrated (1 mg/ml) sterile solution containing PBS, pH 7.4.

Solubility

It is recommended to reconstitute the lyophilized FGF-acidic in sterile 18 MΩ-cm H₂O at 4°C at a concentration of 0.1 mg - 0.25 mg per 1 ml. Allow sample to sit for 5 min. at 4°C, spin to remove precipitant.

Stability

Lyophilized Fibroblast Growth Factor-1, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution FGF-a should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Purity


Greater than 95.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

Amino Acid Sequence

MFNLPPGNYK KPKLLYCSNG GHFLRILPDG TVDGTDRSD QHIQLQLSAE SVGEVYIKST ETGQYLAMDT DGLLYGSQTP
NEECLFLERL EENHYNTYIS KKHAEKNWFV GLKKNNGSCKR GPRTHYGQKA ILFLPLPVSS D

Activity

The ED₅₀, calculated by the dose-dependant proliferation of mouse BALB/c 3T3 cells is <0.5 ng/ml, corresponding to a specific activity of >2.0 x 1,000,000 IU/mg.



Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.