



Interleukin-13, human recombinant (rHuIL-13)

Catalog No: 51562
Lot No: XXXXX
Source: *E. coli*
Synonyms: NC30, ALRH, BHR1, P600, IL-13, MGC116786, MGC116788, MGC116789

Background

IL13 is an immunoregulatory cytokine produced primarily by activated Th2 cells. IL-13 is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen-induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4.

Description

Interleukin-13 human recombinant produced in *E. coli* is a single, non-glycosylated polypeptide chain containing 112 amino acids and having a molecular mass of 12 kDa. IL-13 is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

The protein (1 mg/ml) was lyophilized with 1xPBS pH 7.2 and 5% trehalose.

Solubility

It is recommended to reconstitute the lyophilized Interleukin-13 in sterile 18 M Ω -cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized Interleukin-13, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL-13 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity

Greater than 95% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

Amino Acid Sequence

GPVPPSTALR ELIEELVNIT QNQKAPLCNG SMVWSINLTA GMYCAALES L INVSGCSAIE KTQRMLSGFC PHKVSAGQFS
SLHVRDTKIE VAQFVKDLLL HLKKLFRFREGR FN

Activity

The ED50 was determined by the dose dependent proliferation of TF-1 cells and was found to be <1 ng/ml, corresponding to a specific activity of >1 x 10⁶ units/mg.



Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.