



Anti-GFP (MOUSE) Monoclonal Antibody Biotin Conjugated - 600-306-215

Code: 600-306-215

Size: 1 mg

Product Description: Anti-GFP (MOUSE) Monoclonal Antibody Biotin Conjugated - 600-306-215

Concentration: 1.0 mg/mL by UV absorbance at 280 nm

PhysicalState: Lyophilized

Label	Biotin
Host	Mouse
Species Reactivity	wt, rGFP, eGFP
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Synonyms	mouse anti-GFP antibody biotin conjugation, biotin conjugated mouse anti-GFP antibody, Green Fluorescent Protein, GFP antibody, Green Fluorescent Protein antibody, EGFP, enhanced Green Fluorescent Protein, Aequorea victoria, Jellyfish
Application Note	Monoclonal anti-GFP is designed to detect enhanced GFP and GFP containing recombinant proteins. This antibody can be used to detect GFP by ELISA (sandwich or capture) for the direct binding of antigen. Biotin conjugated monoclonal anti-GFP is well suited to titrate GFP in a sandwich ELISA in combination with Rockland's polyclonal anti-GFP (600-101-215) as the capture antibody. Only use the monoclonal form for the detection of enhanced or recombinant GFP. Polyclonal anti-GFP detects all variants of GFP tested to date. The biotin conjugated detection antibody is typically used with streptavidin conjugated HRP (code # S000-03) or other streptavidin conjugates. The use of polyclonal anti-GFP results in significant amplification of signal when fluorochrome conjugated polyclonal anti-GFP is used relative to the fluorescence of GFP alone. For immunoblotting use either alkaline phosphatase or peroxidase conjugated anti-GFP to detect GFP or GFP containing proteins on western blots. Optimal titers for applications should be determined by the researcher.
Background	Green fluorescent protein is a 27 kDa protein produced from the jellyfish <i>Aequorea victoria</i> , which emits green light (emission peak at a wavelength of 509nm) when excited by blue light. GFP is an important tool in cell biology research. GFP is widely used enabling researchers to visualize and localize GFP-tagged proteins within living cells without the need for chemical staining.
Purity And Specificity	GFP Biotin Conjugated Antibody was prepared from tissue culture supernatant by Protein A affinity chromatography. Assay by Immunoelectrophoresis resulted in a single precipitin arc against anti-biotin and anti-Mouse Serum. Reactivity is observed against recombinant Green Fluorescent Protein (000-001-215, recombinant GFP from <i>Aequorea victoria</i>) by both Western blot and ELISA. No reaction is seen against RFP.
Assay Dilutions	User Optimized
ELISA	1:50,000 - 1:200,000
Western Blot	1:2,000 - 1:10,000
Immunohistochemistry	1:1,000 - 1:5,000
Other Assays	User Optimized
Expiration	Expiration date is one (1) year from date of opening.
Immunogen	Anti-Green Fluorescent Protein (GFP) is produced by immunizing GFP containing fusion protein that corresponds to the full length amino acid sequence (246aa) derived from the jellyfish <i>Aequorea victoria</i> .
Related Products	

200-301-268

Anti-AKT pS473 (MOUSE) Monoclonal Antibody - 200-301-268

610-4302	Anti-MOUSE IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated - 610-4302
611-1302	Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302
B304	NORMAL GOAT SERUM (NGS) - B304

Images


1	Western Blot of Anti-GFP (MOUSE) Monoclonal Antibody Biotin Conjugated antibody. Lane 1: 50ng of GFP. Lane 2: none. Primary antibody: none. Secondary antibody: Anti-GFP (MOUSE) Monoclonal Antibody Biotin Conjugated (p/n 600-306-215) secondary antibody was used at 1:5000 in Blocking Buffer for Fluorescent Western Blotting (p/n MB-070) for 45 min at RT. HRP Streptavidin (p/n S000-03) was used at 1:40,000 in MB-070 for 30 min at 20°C. Block: 5% Blotto (p/n B501-0500) 30 min at 20°C. Predicted/Observed size: 28 kDa for GFP. Other band(s): none.
---	--

1



2

IHC - Immuno-Fluorescence of Biotin Mouse anti-GFP antibody. Biotin mouse anti GFP used 1:5000 As referenced in: Lu S-M, Tremblay M-E, King IL, Qi J, Reynolds HM, et al. (2011) HIV-1 Tat-Induced Microgliosis and Synaptic Damage via Interactions between Peripheral and Central Myeloid Cells. PLoS ONE 6(9): e23915. doi:10.1371/journal.pone.0023915



3

Western Blot of Anti-GFP (MOUSE) Monoclonal Antibody Biotin Conjugated antibody. Lane 1: 50ng of GFP. Lane 2: none. Primary antibody: none. Secondary antibody: Anti-GFP (MOUSE) Monoclonal Antibody Biotin Conjugated (p/n 600-306-215) secondary antibody was used at 1:5000 in Blocking Buffer for Fluorescent Western Blotting (p/n MB-070) for 45 min at RT. HRP Streptavidin (p/n S000-03) was used at 1:40,000 in MB-070 for 30 min at 20°C. Block: 5% Blotto (p/n B501-0500) 30 min at 20°C. Predicted/Observed size: 28 kDa for GFP. Other band(s): none.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.