

Anti-Collagen Type III (RABBIT) Antibody - 600-401-105-0.5

Code: 600-401-105-0.5

Size: 500 µg

Product Description: Anti-Collagen Type III (RABBIT) Antibody - 600-401-105-0.5

Concentration: 1.0 mg/mL by UV absorbance at 280 nm

PhysicalState: Liquid (sterile filtered)

Label	Unconjugated
Host	Rabbit
Gene Name	COL3A1
Species Reactivity	human, bovine
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Volume	500 µL
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at 4° C prior to opening. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage, mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
Synonyms	rabbit anti-Collagen Type III antibody, Collagen type III alpha 1 antibody, Collagen type III alpha antibody, EDS4A antibody, Ehlers Danlos syndrome type IV, autosomal dominant antibody, Fetal collagen antibody, COL3A1, Collagen alpha-1 (III) chain
Application Note	Anti-Collagen antibodies have been used for indirect trapping ELISA for quantitation of antigen in serum using a standard curve, immunoprecipitation, immunohistochemistry, native (non-denaturing, non-dissociating) PAGE, and western blotting for highly sensitive qualitative analysis.
Background	Rockland produces highly active antibodies and conjugates to collagens. Collagens are highly conserved throughout evolution and are characterized by an uninterrupted "Glycine-X-Y" triplet repeat that is a necessary part of the triple helical structure. For these reasons, it is often extremely difficult to generate antibodies with specificities to collagens. The development of 'type' specific antibodies is dependent on NON-DENATURED three-dimensional epitopes. Rockland extensively purifies collagens for immunization from human and bovine placenta and cartilage by limited pepsin digestion and selective salt precipitation. This preparation results in a native conformation of the protein. Antibodies are isolated from rabbit antiserum and are extensively cross-adsorbed by immunoaffinity purification to produce 'type' specific antibodies. Greatly diminished reactivity and selectivity of these antibodies will result if denaturing and reducing conditions are used for SDS-PAGE and immunoblotting.
Purity And Specificity	Anti-Collagen Type III has been prepared by immunoaffinity chromatography using immobilized antigens followed by extensive cross-adsorption against other collagens, human serum proteins and non-collagen extracellular matrix proteins to remove any unwanted specificities. Typically negligible cross-reactivity against other types of collagens was detected by ELISA against purified standards. Some class-specific anti-collagens may be specific for three-dimensional epitopes which may result in diminished reactivity with denatured collagen or formalin-fixed, paraffin embedded tissues. This antibody reacts with most mammalian Type III collagens and has negligible cross-reactivity with Type I, II, IV, V or VI collagens. Non-specific cross-reaction of anti-collagen antibodies with other human serum proteins or non-collagen extracellular matrix proteins is negligible.
Assay Dilutions	User Optimized
ELISA	1:5,000 - 1:50,000
Western Blot	1:1,000 - 1:10,000
Immunohistochemistry	1:50 - 1:200
Other Assays	User Optimized
Expiration	Expiration date is one (1) year from date of opening.
Immunogen	Collagen Type III from human and bovine placenta

Specific Reference

Mak KM, Kwong AJ, Chu E, Hoo NM. (2011) Hepatic Steatosis, Fibrosis, and Cancer in Elderly Cadavers. *Cancer Biology*. 5 DEC 2011. DOI: 10.1002/ar.21525.

Santanu Chakraborty, Elaine E. Wirrig, Robert B. Hinton, Walter H. Merrill, Douglas B. Spicer, Katherine E. Yutzey. (2010) Twist1 promotes heart valve cell proliferation and extracellular matrix gene expression during development in vivo and is expressed in human diseased aortic valves. <https://doi.org/10.1016/j.ydbio.2010.08.021>.

Related Products

001-001-105	Bovine COLLAGEN Type III - 001-001-105
009-001-105	Human COLLAGEN Type III - 009-001-105
600-401-105-0.1	Anti-Collagen Type III (RABBIT) Antibody - 600-401-105-0.1
600-406-105	Anti-Collagen Type III (RABBIT) Antibody Biotin Conjugated - 600-406-105

Related Links

UniProtKB - P02461

<http://www.uniprot.org/uniprot/P02461>


NCBI - P02461.4 <http://www.ncbi.nlm.nih.gov/protein/P02461.4>

GeneID - 1281

Images


1

Rockland anti collagen III antibody (600-401-105 Lot 26016, 1:400, 45 min RT) showed strong staining in FFPE sections of human skin(left, dermis) with moderate to strong red staining and testis (right) where strong staining was observed within connective tissue between seminiferous tubules. The antibody showed strong extracellular staining within connective tissues across many organs with minimal background staining. Slides were steamed in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes for antigen retrieval. Images provided courtesy of LifeSpan Biosciences, Seattle, WA




2

Western Blot of Rabbit Anti-COLLAGEN III Antibody Lane 1:
 Human Collagen III Load: 100 ng per lane
 Primary antibody:
 Collagen III Antibody at 1:1000 o/n at 4°C
 Secondary antibody:
 DyLight™ 649 Goat anti-rabbit at 1:20,000 for 30 min at RT
 Block:
 MB-070 for 30 min at RT
 Predicted/Observed size: 138 kDa, 138 kDa



3

Immunohistochemistry of Rabbit Anti-collagen type III antibody.
 Tissue: right lobe of the liver section. A: Central Vein (CV) fibrosis, B: Non-fibrotic CV, C: Perisinusoidal fibrosis, D: Non-fibrotic area, E: Protat tract fibrosis, F: Septal fibrosis (arrow).
 Fixation: formalin fixed paraaffin embedded. Antigen retrieval: not required.
 Primary antibody: Anti-collagen type III at 1:500 for 4°C for 24hr.
 Secondary antibody: Peroxidase biotin-streptavidin rabbit secondary antibody at 1:10,000 for 45 min at RT.
 Localization: Anti-collagen type III is intra and extracellular.
 Staining: 3,3'-diaminobenzidine tetrahydrochloride was used as the chromogen. Nuclei were counterstained purple with hematoxylin.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.