



Anti-Collagen Type VI (RABBIT) Antibody - 600-401-108-0.1

Code: 600-401-108-0.1

Size: 100 µg

Product Description: Anti-Collagen Type VI (RABBIT) Antibody - 600-401-108-0.1

Concentration: 1 mg/ml by UV absorbance at 280 nm

PhysicalState: Liquid (sterile filtered)

Label	Unconjugated
Host	Rabbit
Gene Name	COL6A1
Species Reactivity	human, bovine
Buffer	0.125 M Sodium Borate, 0.075 M Sodium Chloride, 0.005 M EDTA, pH 8.0
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at 4° C prior to opening. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage, mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
Synonyms	rabbit anti-Collagen Type VI antibody, Collagen alpha-1 (VI) chain, Collagen VI antibody, Human mRNA for collagen VI alpha 1 C terminal globular domain antibody
Application Note	Anti-Collagen antibodies have been used for indirect trapping ELISA for quantitation of antigen in serum using a standard curve, immunoprecipitation, native (non-denaturing, non-dissociating) PAGE, immunohistochemistry, and western blotting for highly sensitive qualitative analysis.
Background	Rockland produces highly active antibodies and conjugates to collagens. Collagens are highly conserved throughout evolution and are characterized by an uninterrupted "Glycine-X-Y" triplet repeat that is a necessary part of the triple helical structure. For these reasons, it is often extremely difficult to generate antibodies with specificities to collagens. The development of 'type' specific antibodies is dependent on NON-DENATURED three-dimensional epitopes. Rockland extensively purifies collagens for immunization from human and bovine placenta and cartilage by limited pepsin digestion and selective salt precipitation. This preparation results in a native conformation of the protein. Antibodies are isolated from rabbit antiserum and are extensively cross-adsorbed by immunoaffinity purification to produce 'type' specific antibodies. Greatly diminished reactivity and selectivity of these antibodies will result if denaturing and reducing conditions are used for SDS-PAGE and immunoblotting.
Purity And Specificity	Anti-Collagen Type VI has been prepared by immunoaffinity chromatography using immobilized antigens followed by extensive cross-adsorption against other collagens, human serum proteins and non-collagen extracellular matrix proteins to remove any unwanted specificities. Typically negligible cross reactivity against other types of collagens was detected by ELISA against purified standards. Some class specific anti-collagens may be specific for three-dimensional epitopes which may result in diminished reactivity with denatured collagen or formalin-fixed, paraffin embedded tissues. This antibody reacts with most mammalian Type VI collagens and has negligible cross-reactivity with Type I, II, III, IV or V collagens. Non-specific cross-reaction of anti-collagen antibodies with other human serum proteins or non-collagen extracellular matrix proteins is negligible.
Assay Dilutions	User Optimized
ELISA	1:5,000 - 1:20,000
Western Blot	1:500 - 1:5,000
Immunohistochemistry	1:50 - 1:200
Other Assays	User Optimized
Expiration	Expiration date is one (1) year from date of opening.
Immunogen	Collagen Type VI from human and bovine placenta

Related Products

001-001-108	Bovine COLLAGEN Type VI - 001-001-108
009-001-108	Human COLLAGEN Type VI - 009-001-108

600-401-108-0.5 Anti-Collagen Type VI (RABBIT) Antibody - 600-401-108-0.5
600-406-108 Anti-Collagen Type VI (RABBIT) Antibody Biotin Conjugated - 600-406-108

Related Links

NCBI - AAH52575.1


<http://www.ncbi.nlm.nih.gov/protein/AAH52575.1>

UniProtKB - <http://www.uniprot.org/uniprot/P12109>

GeneID - 1291

Images

1 Rockland anti collagen VI antibody (600-401-108 Lot 26009, 1:400 45 min RT) showed strong staining in FFPE sections of human placenta (Left) with red staining of stromal and extracellular spaces, and in testis (Right) with staining of extracellular spaces between seminiferous tubules). Slides were steamed in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes for antigen retrieval. Images provided courtesy of LifeSpan Biosciences, Seattle, WA



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.