

Anti-mTOR pS2448 (RABBIT) Antibody - 600-401-422

Code: 600-401-422

Size: 100 µg

Product Description: Anti-mTOR pS2448 (RABBIT) Antibody - 600-401-422

Concentration: 1.1mg/mL by UV absorbance at 280 nm

PhysicalState: Liquid (sterile filtered)

Label	Unconjugated
Host	Rabbit
Gene Name	MTOR
Species Reactivity	human
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Synonyms	rabbit anti-mTOR pS2448 antibody, FKBP12 rapamycin complex associated protein antibody, Serine/threonine-protein kinase mTOR, FK506-binding protein 12-rapamycin complex-associated protein 1, Mammalian target of rapamycin, mTOR, Mechanistic target of rapamycin, Rapamycin and FKBP12 target 1, Rapamycin target protein 1, FRAP, FRAP1, FRAP2, RAFT1, RAPT1
Application Note	This affinity purified antibody has been tested for use in immunohistochemistry, ELISA and western blotting. Western blotting shows reactivity specific for phospho mTOR detecting a band at approximately 250 kDa. Reactivity in other immunoassays is unknown.
Background	Mammalian target of rapamycin (mTOR) is a serine and threonine protein kinase that regulates numerous cellular functions, in particular, the initiation of protein translation. Rapamycin is a natural product macrolide that induces G ₁ growth arrest in yeast, Drosophila, and mammalian cells. mTOR has a long list of synonyms including FK506 binding protein12 - rapamycin associated protein 1, FK506 binding protein12 - rapamycin associated protein 2, FRAP1, FRAP2, RAFT1, RAPT1 and/or FKBP12-rapamycin associated protein (FRAP). mTOR is one of a family of proteins involved in cell cycle progression, DNA recombination, and DNA damage detection. In rat, mTOR is a 245-kD protein referred to as RAFT1 with significant homology to the Saccharomyces cerevisiae protein TOR1 and has been shown to associate with the immunophilin FKBP12 in a rapamycin-dependent fashion. The FKBP12-rapamycin complex is known to inhibit progression through the G ₁ cell cycle stage by interfering with mitogenic signaling pathways involved in G ₁ progression in several cell types, as well as in yeast. The binding of mTOR to FKBP12-rapamycin correlates with the ability of these ligands to inhibit cell cycle progression.
Purity And Specificity	This is an affinity purified antibody produced by immunoaffinity chromatography using the immunizing peptide after immobilization to a solid phase. Reactivity occurs with phosphorylated mTOR from human derived tissues and cells. Reactivity against mTOR from other species has not been determined, however, reactivity with mouse and rat is suggested based on protein sequence homologies.
Assay Dilutions	User Optimized
ELISA	1:10,000 - 1:100,000
Western Blot	1:500 - 1:2,000
Immunohistochemistry	5.0 µg/ml
Other Assays	User Optimized
Expiration	Expiration date is one (1) year from date of opening.
Immunogen	Anti-mTOR pS2448 antibody was prepared from whole rabbit serum produced by repeated immunizations with a phosphorylated synthetic peptide corresponding a c-terminal region near Serine 2448 of human mTOR.

General Reference

Kristof,A.S. et al. (2003) Stimulation of signal transducer and activator of transcription-1 (STAT1)-dependent gene transcription by lipopolysaccharide and interferon-gamma is regulated by mammalian target of rapamycin. *J. Biol. Chem.* 278 (36), 33637-33644.

Chen,Y., et al. (2003) Phospholipase D confers rapamycin resistance in human breast cancer cells. *Oncogene* 22 (25), 3937-3942.

Nojima,H. et al. (2003) The mammalian target of rapamycin (mTOR) partner, raptor, binds the mTOR substrates p70 S6 kinase and 4E-BP1 through their TOR signaling (TOS) motif. *J. Biol. Chem.* 278 (18), 15461-15464.

Related Products

100-401-401	Anti-AKT (RABBIT) Antibody - 100-401-401
200-301-268	Anti-AKT pS473 (MOUSE) Monoclonal Antibody - 200-301-268
200-301-269	Anti-AKT pT308 (MOUSE) Monoclonal Antibody - 200-301-269
600-401-268	Anti-AKT pS473 (RABBIT) Antibody - 600-401-268

Related Links

NCBI - 1169735


<http://www.ncbi.nlm.nih.gov/protein/1169735>

UniProtKB - <http://www.uniprot.org/uniprot/P42345>


GeneID - 2475

Images


- 1 Rockland's affinity purified anti-mTOR pS 2448 antibody was used at 5µg/ml to detect signal in a variety of tissues including multi-human, multi-brain and multi-cancer slides. This image shows moderate staining of proximal convoluted tubules of the kidney. Tissue was formalin-fixed and paraffin embedded. The image shows localization of the antibody as the precipitated red signal, with a hematoxylin purple nuclear counterstain. Personal Communication, Tina Roush, LifeSpanBiosciences, Seattle, WA.



- 2 Affinity Purified Anti-mTOR pS 2448 (Rabbit) is shown to detect a 250 kDa band (indicated) corresponding to phosphorylated human mTOR present in a 293T whole cell lysates. Cells were serum-starved for 24 hours prior to harvest. ~20 µg of lysate was loaded per lane for SDS-PAGE. Untreated cells are shown in lane 1, whereas cells in lane 2 were treated with IGF-1 (100 ng/ml) for 20 min prior to harvest. Follow reaction of antibody with a 1:2000 dilution of HRP Goat-a-Rabbit IgG for visualization.



3 Metabolic and autocrine regulation of the mTOR pathway by Beta-cells.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.