

Anti-ABCB5 (RABBIT) Antibody - 600-401-A77S

Code: 600-401-A77S

Size: 25 µL

Product Description: Anti-ABCB5 (RABBIT) Antibody - 600-401-A77S

Concentration: 0.31 mg/mL by UV absorbance at 280 nm

PhysicalState: Liquid (sterile filtered)

Label	Unconjugated
Host	Rabbit
Gene Name	ABCB5
Species Reactivity	human
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Synonyms	rabbit anti-ABCB5 Antibody, ABCB 5, ABCB-5, ATP binding cassette sub family B (MDR/TAP) member 5 antibody, ATP binding cassette sub family B member 5 antibody, EST422562 antibody, P glycoprotein ABCB5 antibody
Application Note	This affinity purified antibody has been tested for use in ELISA, IHC, and western blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 117 kDa in size corresponding to ABCB5 by western blotting in the appropriate cell lysate or extract.
Background	This antibody is designed, produced, and validated as part of a collaboration between Rockland and the National Cancer Institute (NCI) and is suitable for Cancer, Immunology and Nuclear Signaling research. Anti-ABCB5 Antibody is a ATP-Binding Cassette protein. ABCB5 is a novel member of the human P-glycoprotein family. It functions as a determinant of membrane potential and regulator of cell fusion in physiologic skin progenitor cells. Cell fusion is thought to contribute to tissue differentiation. ABCB5 is predominantly expressed in melanoma cells and is a novel molecular marker for a distinct subset of chemoresistant stem cell phenotype-expressing tumor cells among human epidermal melanocytes. ABCB5 is a rhodamine efflux transporter. This antibody product is intended to be used to confirm cellular localization and expression level of ABCB5. ABCB5 antibody is ideal for Cancer and Signal Transduction research.
Purity And Specificity	This product was affinity purified from monospecific antiserum by immunoaffinity chromatography. This antibody is specific for human ABCB5 protein. A BLAST analysis was used to suggest partial cross-reactivity with ABCB5 from monkey (85% homology), rat (68% homology) and mouse (62% homology) sources. Cross-reactivity with ABCB5 from other sources has not been determined.
Assay Dilutions	User Optimized
ELISA	1:10,000 - 1:15,000
Western Blot	1:10,000
Immunohistochemistry	1:200
Other Assays	User Optimized
Expiration	Expiration date is three (3) months from date of opening.
Immunogen	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to residues corresponding to an internal region of human ABCB5.
General Reference	Frank NY, Margaryan A, Huang Y, Schatton T, Waaga-Gasser AM, Gasser M, Sayegh MH, Sadee W, Frank MH. (2005) ABCB5-mediated doxorubicin transport and chemoresistance in human malignant melanoma. <i>Cancer Res.</i> 65(10):4320-33. Chen KG, Szakacs G, Annereau JP, Rouzaud F, Liang XJ, Valencia JC, Nagineni CN, Hooks JJ, Hearing VJ, Gottesman MM. (2005) Principal expression of two mRNA isoforms (ABCB 5alpha and ABCB 5beta) of the ATP-binding cassette transporter gene ABCB 5 in melanoma cells and melanocytes. <i>Pigment Cell Res.</i> 18(2):102-12. Chen KG, Gottesman MM. (2005) Useful tool to generate unidirectional deletion vectors by utilizing the star activity of BamHI in an NcoI-BamHI-XhoI cassette. <i>Biotechniques.</i> 38(2):198, 200, 202, 204.

Related Products

009-001-B99	VEGF-165 Human Recombinant Protein - 009-001-B99
009-001-C23	VEGF-121 Human Recombinant Protein - 009-001-C23
200-401-B74	Anti-Bovine VEGF-A (RABBIT) Antibody - 200-401-B74
200-401-B74S	Anti-Bovine VEGF-A (RABBIT) Antibody - 200-401-B74S

Related Links

UniProtKB - Q2M3G0

<http://www.uniprot.org/uniprot/Q2M3G0>


NCBI - 56849536 <http://www.ncbi.nlm.nih.gov/protein/56849536>

GenID - 340273

Images


1

Immunohistochemistry of ABCB5 antibody Tissue: human liver
Fixation: formalin fixed paraffin embedded Antigen retrieval: user optimized Primary antibody: ABCB5 1:200 Secondary antibody: Peroxidase goat anti-rabbit at 1:10,000 for 45 min at RT
Localization: Moderate to strong cytoplasmic and membranous staining was observed in hepatocytes. Occasional nuclear staining was observed in hepatocytes and sinusoidal cells. Staining: antibody as precipitated red signal with a hematoxylin purple nuclear counterstain






2

Western blot using Rockland's affinity purified anti-ABCB5 antibody shows detection of ABCB5 beta in ~12.5 ug of transfected-Hi5 whole cell lysate. No reaction was seen when antibody was pre-incubated with the immunizing peptide (data not shown). A 3-8% Tris-acetate gel was used for separation. The arrowhead corresponds to 117 kDa ABCB5. The membrane was probed with the primary antibody at a 1:10,000 dilution in 5% milk in TBST at 4° C, overnight. Personal Communication, JP Gillet, CCR-NCI, Bethesda, MD.





3

Immunohistochemistry of Rabbit anti ABCB5 antibody. Tissue: human breast carcinoma at pH6. Fixation: formalin fixed paraffin embedded. Primary antibody: ABCB5 antibody at 10 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: ABCB5 is cytoplasmic. Staining: ABCB5 as precipitated brown signal with hematoxylin purple nuclear counterstain.



4

Immunohistochemistry of Rabbit anti ABCB5 antibody. Tissue: human breast carcinoma at pH9. Fixation: formalin fixed paraffin embedded. Primary antibody: ABCB5 antibody at 10 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: ABCB5 is cytoplasmic. Staining: ABCB5 as precipitated brown signal.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.