

Anti-DISC1 (RABBIT) Antibody - 600-401-AV9

Code: 600-401-AV9

Size: 100 µg

Product Description: Anti-DISC1 (RABBIT) Antibody - 600-401-AV9

Concentration: 1 mg/mL by UV absorbance at 280 nm

PhysicalState: Liquid (sterile filtered)

Label	Unconjugated
Host	Rabbit
Gene Name	DISC1
Species Reactivity	Human, mouse
Buffer	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Stabilizer	None
Preservative	0.02% (w/v) Sodium Azide
Storage Condition	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Synonyms	DISC1 Antibody, SCZD9, C1orf136, KIAA0457, Disrupted in schizophrenia 1 protein
Application Note	Anti-DISC1 Antibody has been tested for use in ELISA, Western Blotting, Immunocytochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 94 kDa in Western Blots of specific cell lysates and tissues.
Background	Disrupted in schizophrenia 1 (DISC1) is a candidate gene for susceptibility to schizophrenia. It was discovered through chromosomal analysis of a large Scottish family whose members exhibited schizophrenia and related psychiatric disorders. Through yeast two-hybrid screening, it was discovered that DISC1 interacts with many members of the centrosome and cytoskeletal system including MAP1A and Nudel. More recently, DISC1 has been found to regulate the transport of a complex containing Nudel, the lissencephaly-1 (LIS1) protein, and 14-3-3epsilon from neuronal cell bodies to the axons by the action of the microtubule-dependent directed motor protein kinesin-1, also known as KIF5A. Decreased expression of DISC1 in neurons caused an accelerated rate of neuronal integration, resulting in aberrant morphological development, suggesting that DISC1 plays a role in dendritic development and synapse formation. DISC1 has at least four known isoforms.
Purity And Specificity	Anti-DISC1 Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. Cross reactivity with DISC1 from other sources has not been determined.
ELISA	1:5000-1:20,000
Western Blot	0.5-2 g/mL
Immunohistochemistry	5 g/mL
IF Microscopy	20 g/mL
Expiration	Expiration date is one (1) year from date of opening.
Immunogen	Anti-DISC1 antibody was prepared from whole rabbit serum produced by repeated immunizations with a 18 amino acid synthetic peptide from near the internal region of human DISC1.

Related Products

200-301-401	Anti-AKT (MOUSE) Monoclonal Antibody - 200-301-401
611-1302	Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302
BSA-50	BOVINE SERUM ALBUMIN - Fraction V (Immunoglobulin and Protease Free) - BSA-50
MB-070	Blocking Buffer for Fluorescent Western Blotting - MB-070

Related Links

UniProtKB - Q9NRI5


<http://www.uniprot.org/uniprot/Q9NRI5>

GeneID - 27185 <http://www.ncbi.nlm.nih.gov/gene/27185>


NCBI - NP_061132

Images

- 1 Immunofluorescence Microscopy of DISC1 antibody. Tissue: HeLa cells. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: DISC1 antibody at 20 $\mu\text{g}/\text{mL}$ for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT. Staining: DISC1 as a red fluorescent signal.




- 2 Immunocytochemistry of DISC1 antibody. Tissue: HeLa cells. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: DISC1 antibody at 5 $\mu\text{g}/\text{mL}$ for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: DISC1 is nuclear and cytoplasmic. Staining: DISC1 as precipitated brown signal with hematoxylin purple counterstain.



3

Western Blot of DISC1 antibody. Lane A: SK-N-SH cell lysate at 0.5 µg/mL. Lane B: SK-N-SH cell lysate at 1 µg/mL. Lane C: SK-N-SH cell lysate at 2 µg/mL. Load: 35 µg per lane. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 93.6 kDa, ~93 kDa for DISC1. Other band(s): DISC1 splice variants and isoforms.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.