

Anti-RUNX1 (RABBIT) Antibody - 600-401-EC9
Code: 600-401-EC9

Size: 100 µg

Product Description: Anti-RUNX1 (RABBIT) Antibody - 600-401-EC9

Concentration: 1 mg/mL by UV absorbance at 280 nm

PhysicalState: Liquid (sterile filtered)

Label	Unconjugated
Host	Rabbit
Gene Name	RUNX1
Species Reactivity	Human, mouse, rat
Buffer	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Stabilizer	None
Preservative	0.02% (w/v) Sodium Azide
Storage Condition	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Synonyms	RUNX1 Antibody, AML1, CBFA2, EVI-1, AMLCR1, PEBP2aB, AML1-EVI-1, AML1, Runt-related transcription factor 1, Acute myeloid leukemia 1 protein, CBF-alpha-2
Application Note	Anti-RUNX1 Antibody has been tested for use in ELISA and Western Blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 49 kDa in Western Blots of specific cell lysates and tissues.
Background	RUNX1 is one of three mammalian RUNX genes that control multiple aspects of embryonic development and are responsible for the pathogenesis of many human diseases. RUNX1 plays major roles in the development of nociceptive sensory neurons in addition to hematopoietic stem cells (HSC) with the exception of the erythroid lineage. During development, Notch signals mediate RUNX1 induction with SCL/GATA/Ets factors, and Wnt signals potentially cooperate with RUNX1 to facilitate adult HSC expansion via cooperative induction of cyclin D, cdk4, and other cell cycle regulators. In turn, RUNX1 regulates cell cycle transitions dependent on functional/physical interactions with other proteins such as HDAC1 and -3, mSin3A, p300, SMAD proteins, and LEF/TCF.
Purity And Specificity	Anti-RUNX1 Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. Cross reactivity with RUNX1 from other sources has not been determined.
ELISA	1:10,000-1:20,000
Western Blot	1-2 g/mL
Expiration	Expiration date is one (1) year from date of opening.
Immunogen	Anti-RUNX1 antibody was prepared from whole rabbit serum produced by repeated immunizations with a 16 amino acid synthetic peptide from near the internal region of human RUNX1.

Related Products

200-301-401	Anti-AKT (MOUSE) Monoclonal Antibody - 200-301-401
611-1302	Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302
BSA-50	BOVINE SERUM ALBUMIN - Fraction V (Immunoglobulin and Protease Free) - BSA-50
MB-070	Blocking Buffer for Fluorescent Western Blotting - MB-070

Related Links

UniProtKB - Q01196

<http://www.uniprot.org/uniprot/Q01196>

UniProtKB -
P59594

<http://www.uniprot.org/uniprot/P59594>

GeneID - 861

<http://www.ncbi.nlm.nih.gov/gene/861>

GeneID - 1489668

<http://www.ncbi.nlm.nih.gov/gene/1489668>

NCBI - AAI36381

<http://www.ncbi.nlm.nih.gov/protein/AI36381>


NCBI - P59594

<http://www.ncbi.nlm.nih.gov/protein/P59594>

Images

1

Western Blot of RUNX1 antibody. Lane 1: Raji cell lysate with RUNX1 antibody at 1 µg/mL. Lane 2: Raji cell lysate with RUNX1 antibody at 2 µg/mL. Load: 35 µg per lane. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 49 kDa, 55 kDa for RUNX1. Other band(s): RUNX1 splice variants and isoforms.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.