



Anti-GOLDEN SYRIAN HAMSTER IgG (H&L) (RABBIT) Antibody Biotin Conjugated - 607-4602

Code: 607-4602

Size: 1.5 mg

Product Description: Anti-GOLDEN SYRIAN HAMSTER IgG (H&L) (RABBIT) Antibody Biotin Conjugated - 607-4602

Concentration: 1.5 mg/mL by UV absorbance at 280 nm

PhysicalState: Lyophilized

Label	Biotin
Host	Rabbit
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Synonyms	Golden Syrian Hamster IgG (H&L) Antibody, Rabbit-a-Golden Syrian Hamster IgG, Golden Syrian Hamster IgG in Rabbit, Rabbit IgG Secondary Antibody, rabbit anti-hamster IgG antibody
Application Note	Anti-Golden Syrian Hamster IgG has been assayed against 1.0 ug of Hamster IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:15,000 to 1:60,000 of the reconstitution concentration is suggested for Anti-Golden Syrian Hamster IgG.
Background	Anti-Golden Syrian Hamster IgG Biotin Antibody generated in rabbit detects Golden Syrian Hamster IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. Anti-Golden Syrian Hamster IgG Antibody is ideal for investigators in Cancer, Immunology, and Microbiology research.
Purity And Specificity	Anti-Golden Syrian Hamster IgG Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Hamster IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Hamster IgG and Hamster Serum.
Assay Dilutions	User Optimized
ELISA	1:20,000 - 1:100,000
Western Blot	1:2,000 - 1:10,000
Immunohistochemistry	1:1,000 - 1:5,000
Other Assays	User Optimized
Expiration	Expiration date is one (1) year from date of opening.
Immunogen	Hamster IgG whole molecule
General Reference	Janeway, Jr., Travers, Walport, and Shlomchik. "The Immune System in Health and Disease." Immunobiology, 5th Edition: Garland Science: 2001.
Related Products	

007-0102

HAMSTER IgG whole molecule - 007-0102

007-0204	HAMSTER IgG F(ab') ₂ fragment Fluorescein conjugated - 007-0204
107-1102	Anti-GOLDEN SYRIAN HAMSTER IgG (H&L) (GOAT) Antibody - 107-1102
107-4101	Anti-GOLDEN SYRIAN HAMSTER SERUM (RABBIT) Antibody - 107-4101

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.