



F(ab')₂ Anti-MOUSE IgG F(ab')₂ (GOAT) Antibody Phycoerythrin conjugated Min X Bv Hs & Hu Serum Proteins - 710-1831

Code: 710-1831

Size: 500 µg

Product Description: F(ab')₂ Anti-MOUSE IgG F(ab')₂ (GOAT) Antibody Phycoerythrin conjugated Min X Bv Hs & Hu Serum Proteins - 710-1831

Concentration: 0.5 mg/mL by absorbance = 82.0 at 565 nm

PhysicalState: Lyophilized

Label	R-Phycoerythrin (RPE)
Host	Goat
Emission Wavelength	575
Excitation Wavelength	488
Species Reactivity	Mouse
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide
Storage Condition	Store vial at 4° C prior to restoration. Restore with deionized water (or equivalent). This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. Centrifuge product if not completely clear after standing at room temperature. Do not freeze after reconstitution. Store reagent in the dark. Use subdued lighting during handling and incubation of cells prior to analysis.
Synonyms	Goat F(ab') ₂ Anti-Mouse IgG F(ab') ₂ Antibody phycoerythrin Conjugation, Goat Fab2 Anti-Mouse IgG Fab2 PE Conjugated Antibody
Application Note	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10 ⁶ cells in flow cytometry is approximately 1.0 µg of antibody conjugate. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.
Background	F(ab') ₂ Anti-Mouse IgG F(ab') ₂ Phycoerythrin Antibody generated in goat detects Mouse F(ab') ₂ . Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(ab') ₂ Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.
Purity And Specificity	F(ab') ₂ fragment PE conjugated Anti-Mouse IgG F(ab') ₂ was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Phycoerythrin, anti-Goat Serum, Mouse IgG, Mouse IgG F(ab') ₂ and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Mouse IgG F(c) or Bovine, Horse or Human Serum Proteins.
Assay Dilutions	FLOW 1:100 - 1:250
Western Blot	1:100 - 1:500
IF Microscopy	1:100 - 1:250
Flow Cytometry	1:100 - 1:250
Other Assays	FLOW 1:100 - 1:250
Expiration	Expiration date is six (6) months from date of opening.
Immunogen	Mouse IgG F(ab') ₂ fragment


Related Products

200-301-268	Anti-AKT pS473 (MOUSE) Monoclonal Antibody - 200-301-268
B304	NORMAL GOAT SERUM (NGS) - B304
MB-070	Blocking Buffer for Fluorescent Western Blotting - MB-070
PA00-41	PROTEIN A DyLight™ 488 Conjugated - PA00-41

Images

1

Western blot of Phycoerythrin conjugated Goat F(ab')₂ Anti-Mouse IgG F(ab')₂ Pre-Adsorbed secondary antibody. Lane 1: Mouse Fab2. Lane 2: None. Load: 50 ng per lane. Primary antibody: None. Secondary antibody: Phycoerythrin goat secondary antibody at 1:1,000 for 60 min at RT. Blocking: MB-070 for 30 min at RT. Predicted/Observed size: 25 kDa, 25 kDa for Mouse IgG F(ab')₂. Other band(s): None.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.