



Interleukin-17, mouse recombinant (rmIL-17)

Catalog No: 87321
Lot No: XXXXX
Source: *E. coli*
Synonyms: CTLA-8, IL-17, IL-17A, Cytotoxic T-lymphocyte-associated antigen 8

Background

IL17 is a proinflammatory cytokine produced by activated T cells. IL-17 regulates the activities of NF-kappaB and mitogen-activated protein kinases. Interleukin-17 can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of IL-17 are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis.

Description

Interleukin-17 murine recombinant produced in *E. coli* is a homodimeric, non-glycosylated polypeptide chain containing a total of 268 (2x134) amino acids and having a molecular mass of 30 kDa. IL-17 is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

Lyophilized from a concentrated (1 mg/ml) solution containing no additives.

Solubility

It is recommended to reconstitute the lyophilized Interleukin-17 in sterile 18 MΩ-cm H₂O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized Interleukin-17, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL-17 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity

Greater than 98.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

Amino Acid Sequence

MAAIIPQSSA CPNTEAKDFL QNVKVNKVF NSLGAKVSSR RPSDYLNRST SPWTLHRNED PDRYPSVIWE AQCRHQRCVN
AEGKLDHNMN SVLIQQEILV LKREPESCPF TFRVEKMLVG VGCTCVASIV RQAA

Activity

The ED50 as determined by the dose-dependant induction of IL-6 production in cultured mouse NIH 3T3 fibroblasts was found to be 0.4 ng/ml, corresponding to a specific activity of 2,500,000 units/mg.

CONTACT US TODAY

BIOMOL GmbH • Kieler Straße 303a • 22525 Hamburg • Germany • info@biomol.de • www.biomol.de

Fon: +49 (0)40-853 260 0 • TOLL FREE IN GERMANY: Fon: 0800-246 66 51



Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.

CONTACT US TODAY

BIOMOL GmbH • Kieler Straße 303a • 22525 Hamburg • Germany • info@biomol.de • www.biomol.de
Fon: +49 (0)40-853 260 0 • TOLL FREE IN GERMANY: Fon: 0800-246 66 51