



Insulin-Like Growth Factor-I, rat recombinant (rrIGF-1)

Catalog No: 87394
Lot No: XXXXX
Source: *E. coli*
Synonyms: Somatomedin C, IGF-I, IGFIA, IGF1

Background

The somatomedins, or insulin-like growth factors (IGFs), comprise a family of peptides that play important roles in mammalian growth and development. IGF1 mediates many of the growth-promoting effects of growth hormone (GH; MIM 139250). Early studies showed that growth hormone did not directly stimulate the incorporation of sulfate into cartilage, but rather acted through a serum factor, termed 'sulfation factor,' which later became known as 'somatomedin' (Daughaday et al., 1972). Three main somatomedins have been characterized: somatomedin C (IGF1), somatomedin A (IGF2; MIM 147470), and somatomedin B (MIM 193190) (Rotwein, 1986; Rosenfeld, 2003).

Description

Insulin-Like Growth Factor I rat recombinant produced in *E. coli* is a single, non-glycosylated, polypeptide chain containing 70 amino acids and having a molecular mass of 7687 Dalton. IGF-I is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

The protein was lyophilized with no additives.

Solubility

It is recommended to reconstitute the lyophilized IGF1 in sterile 18 MΩ-cm H₂O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized IGF-1, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IGF1 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity

Greater than 98.0% as determined by SDS-PAGE.

Activity

The ED₅₀, was calculated 2 methods: 1. Stimulation of protein synthesis in rat L6 myoblasts ED₅₀ was found to be <30 ng/ml corresponding to a specific activity of 33,334 IU/mg. 2. Type 1 IGF receptor binding assay ED₅₀ was found to be <10 ng/ml corresponding to a specific activity of 100,000 IU/mg.

Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.