

# Adiponectin, trimeric form, human recombinant (rHuAcrp30-tri)

Catalog No: 94888 Lot No: XXXXX Source: HEK293

**Synonyms:** Acrp30, AdipoQ, GBP-28, APM-1, ACDC

### **Background**

Adiponectin is a hormone exclusively expressed from adipose tissue. Many studies demonstrate that Adiponectin has direct anti-diabetic, anti-atherogenic and anti-inflammatory functions. APM-1 can increase insulin sensitivity of skeletal muscle. Attenuate hepatic lipogenesis and gluconeogenesis, regulate NO production in endothelial cells, inhibit proliferation of smooth muscle cells and prevent lipid accumulation of macrophage cells. In the circulation, Adiponectin is present as three different oligomeric complexes, including the high molecular weight (HMW), the middle molecular weight (MMW, also called hexamer) and low molecular weigh (MMW, also called trimer) forms 8. Different oligomeric complex of Adiponectin activates different signaling pathways and exerts distinct functions.

### Description

Trimeric form of Adiponectin human recombinant was expressed in HEK293 cells. The cysteine 39 was replaced with Alanine (C39A) 9. hAd-C39A can only form a trimer, but not a hexamer or an HMW form.

### **Physical Appearance**

Filtered white lyophilized (freeze-dried) powder.

### **Formulation**

Filtered (0.4  $\mu$ m) and lyophilized from 0.5 mg/ml in 0.05 M phosphate buffer, 0.05 M NaCl, pH 7.2.

## Solubility

Add deionized water to a working volume of 0.5 mg/ml and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

#### Stability

Store lyophilized Adiponectin at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.

### Purity

Greater than 95% as determined by SDS-PAGE.

### Activity

ED50 =  $3 - 8.5 \mu g/ml$ , as determined by its ability to inhibit proliferation of HASMCs induced by HB EGF.

### Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.