



Myostatin, Propeptide, human recombinant (rHuGDF-8)

Catalog No: 94982
Lot No: XXXXX
Source: *E. coli*
Synonyms: GDF-8, MSTN, Growth Differentiation Factor 8, MSTN Muscle Hypertrophy

Background

Myostatin (GDF-8), a member of the TGFbeta superfamily, is a potent and specific negative regulator of skeletal muscle mass. In serum, myostatin circulates as part of a latent complex containing myostatin propeptide and/or follistatin-related gene. The myostatin propeptide is known to bind and inhibit myostatin in vitro. This interaction is relevant in vivo, with a majority (>70%) of myostatin in serum bound to its propeptide. The myostatin propeptide is negative regulator of myostatin in vivo.

Description

Myostatin propeptide human recombinant is a 27.8 kDa protein containing 244 amino acid residues of human Myostatin Propeptide.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

Lyophilized from 10 mM acetic acid.

Solubility

It is recommended to reconstitute the lyophilized Myostatin Propeptide in sterile 50 mM acetic acid at 0.1 mg/ml, which can then be further diluted to other aqueous solutions.

Stability

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time.

Purity

Greater than 95.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

Amino Acid Sequence

MNENSEQKEN VEKEGLCNAC TWRQNTKSSR IEAIIQIILS KLRLETAPNI SKDVIRQLLP KAPPLRELID QYDVQRDSS
DGSLEDDDYH ATTETIITMP TESDFLMQVD GPKKCCFFKF SSKIYQNKVV KAQLWIYLRP VETPTTVFVQ ILRLIKPMKD
GTRYTGIRSL KLDMPGTGI WQSIDVKTVL QNWLKQPESN LGIEIKALDE NGHDLAVTFP GPGEDGLNPF LEVKVTDTPK
RSRR

Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.