



Platelet Factor 4, human recombinant (rHuPF-4)

Catalog No: 97207
Lot No: XXXXX
Source: *E. coli*
Synonyms: CXCL4, PF-4, PF4, Iroplact, Oncostatin-A, SCYB4, MGC138298

Background

Platelet factor-4 is a 70-amino acid protein that is released from the alpha-granules of activated platelets and binds with high affinity to heparin. Its major physiologic role appears to be neutralization of heparin-like molecules on the endothelial surface of blood vessels, thereby inhibiting local antithrombin III activity and promoting coagulation. As a strong chemoattractant for neutrophils and fibroblasts, PF4 probably has a role in inflammation and wound repair. Oncostatin-A is a member of the CXC chemokine family. Human PF4 is used for the proof of heparin-induced thrombocytopenia. Furthermore it is used as an inhibitor in the angiogenesis during tumor therapy.

Description

CXCL4 human recombinant produced in *E. coli* is a single, non-glycosylated polypeptide chain containing 70 amino acids and having a molecular mass of 7.8 kDa.

Physical Appearance

Sterile filtered white lyophilized powder.

Formulation

The CXCL4 protein was lyophilized after extensive dialysis against 50 mM Tris-HCl pH 8.0 and 150 mM NaCl buffer.

Solubility

It is recommended to reconstitute the lyophilized CXCL4 in sterile 18 M Ω -cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized CXCL4, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CXCL4 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity

Greater than 95.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

Amino Acid Sequence

The sequence of the first four N-terminal amino acids was determined and was found to be Glu-Ala-Glu-Glu-Asp.

Activity

The ED₅₀ of CXCL4 as determined by its ability to inhibit human FGF basic dependent proliferation of NR6R3T3 mouse fibroblasts was found to be 5 - 15 μ g/ml.

CONTACT US TODAY



Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.

CONTACT US TODAY

BIOMOL GmbH • Kieler Straße 303a • 22525 Hamburg • Germany • info@biomol.de • www.biomol.de

Fon: +49 (0)40-853 260 0 • TOLL FREE IN GERMANY: Fon: 0800-246 66 51