



## Vascular Endothelial Growth Factor (121 amino acids), mouse recombinant (rmVEGF-121)

**Catalog No:** 97216  
**Lot No:** XXXXX  
**Source:** *E. coli*  
**Synonyms:** Vascular endothelial growth factor A, VEGF-A, Vascular permeability factor, VPF, VEGF, MGC70609

### Background

Vascular endothelial growth factor is an important signaling protein involved in both vasculogenesis and angiogenesis. As its name implies, VEGF activity has been mostly studied on cells of the vascular endothelium, although it does have effects on a number of other cell types (e.g. stimulation monocyte/macrophage migration, neurons, cancer cells, kidney epithelial cells). VEGF mediates increased vascular permeability, induces angiogenesis, vasculogenesis and endothelial cell growth, promotes cell migration, and inhibits apoptosis. In vitro, VEGF has been shown to stimulate endothelial cell mitogenesis and cell migration. VEGF is also a vasodilator and increases microvascular permeability and was originally referred to as vascular permeability factor. Elevated levels of this protein is linked to POEMS syndrome, also known as Crow-Fukase syndrome. Mutations in this gene have been associated with proliferative and nonproliferative diabetic retinopathy.

### Description

Vascular Endothelial Growth Factor-121 mouse recombinant produced in *E. coli* is a homodimer, non-glycosylated, polypeptide chain containing 121 amino acids and having a molecular mass of 28.4 kDa. Recombinant mouse VEGF-121 is a truncated version of murine VEGF-165. VEGF-121 is purified by proprietary chromatographic techniques.

### Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

### Formulation

The protein was lyophilized from a concentrated (1 mg/ml) solution with no additives.

### Solubility

It is recommended to reconstitute the lyophilized VEGF-121 in sterile 18 MΩ-cm H<sub>2</sub>O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

### Stability

Lyophilized VEGF-121, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution VEGF-121 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

### Purity

Greater than 97.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

### Amino Acid Sequence

MAPTTEGEQK SHEVIKFMVDV YQRSYCRPIE TLVDIFQEYYP DEIEYIFKPS CVPLMRCAGC CNDEALECVP TSESNITMQI  
MRIKPHQSQH IGEMSFLQHS RCECRPKKDR TKPEKCDKRPR R

**CONTACT US TODAY**

BIOMOL GmbH • Kieler Straße 303a • 22525 Hamburg • Germany • info@biomol.de • www.biomol.de

Fon: +49 (0)40-853 260 0 • TOLL FREE IN GERMANY: Fon: 0800-246 66 51



### Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.

**CONTACT US TODAY**

BIOMOL GmbH • Kieler Straße 303a • 22525 Hamburg • Germany • [info@biomol.de](mailto:info@biomol.de) • [www.biomol.de](http://www.biomol.de)

Fon: +49 (0)40-853 260 0 • TOLL FREE IN GERMANY: Fon: 0800-246 66 51