



Interleukin-5, mouse recombinant (rmlL-5)

Catalog No: 97304
Lot No: XXXXX
Source: *E. coli*
Synonyms: EDF, BCDFII, TRF, T-cell replacing factor, Eosinophil differentiation factor, B cell differentiation factor II, IL-5, BCGF-II, Cytotoxic T-lymphocyte inducer, IL5

Background

The protein encoded by this gene is a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. This cytokine is a main regulator of eosinopoiesis, eosinophil maturation and activation. The elevated production of this cytokine is reported to be related to asthma or hypereosinophilic syndromes. The receptor of this cytokine is a heterodimer, whose beta subunit is shared with the receptors for interleukin 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene, together with those for interleukin 4 (IL4), interleukin 13 (IL13), and CSF2, form a cytokine gene cluster on chromosome 5. This cytokine, IL4, and IL13 are found to be regulated coordinately by long-range regulatory elements spread over 120 kilobases on chromosome 5q31.

Description

Interleukin-5 mouse recombinant produced in *E. coli* is a dimeric, non-glycosylated polypeptide chain containing 2 x 113 amino acids forming a disulfide linked homodimer and having a molecular mass of 26.2 kDa. IL-5 is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

Lyophilized from a concentrated (1 mg/ml) solution containing 20 mM Sodium Phosphate and 50 mM NaCl pH 7.5.

Solubility

It is recommended to reconstitute the lyophilized Interleukin-5 in sterile 18 M Ω -cm H₂O at a concentration of 100 μ g/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized Interleukin-5, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution mouse IL-5 should be stored at 4°C between 2-7 days and for future use below -18°C. Please avoid freeze thaw cycles.

Purity

Greater than 97.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

Amino Acid Sequence

MEIPMSTVVK ETLTQLSAHR ALLTSNETMR LPVPTHKNHQ LCIGEIFQGL DILKNQTVRG GTVEMLFQNL SLIKKYIDRQ
KEKCGEERRR TRQFLDYLQE FLGVMSTEWA MEG

Activity

The ED50 as determined by the dose-dependant stimulation of the proliferation of TF-1 cells was found to be 1.7 ng/ml, corresponding to a specific activity of 588,235.3 IU/mg.

CONTACT US TODAY



Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.

CONTACT US TODAY

BIOMOL GmbH • Kieler Straße 303a • 22525 Hamburg • Germany • info@biomol.de • www.biomol.de

Fon: +49 (0)40-853 260 0 • TOLL FREE IN GERMANY: Fon: 0800-246 66 51