



Granulocyte Macrophage-Colony Stimulating Factor, rhesus macaque recombinant (rrmGMCSFF)

Catalog No: 97365
Lot No: XXXXX
Source: *E. coli*
Synonyms: CSF-2, MGI-1GM, GM-CSF, Pluripoietin-alpha, Molgramostin, Sargramostim, MGC131935, MGC138897

Background

GMCSF is a cytokine that controls the production, differentiation, and function of granulocytes and macrophages. The active form of the protein is found extracellularly as a homodimer. This gene has been localized to a cluster of related genes at chromosome region 5q31, which is known to be associated with interstitial deletions in the 5q- syndrome and acute myelogenous leukemia. Other genes in the cluster include those encoding interleukins 4, 5, and 13. GM-CSF stimulates the growth and differentiation of hematopoietic precursor cells from various lineages, including granulocytes, macrophages, eosinophils and erythrocytes.

Description

Granulocyte Macrophage Colony Stimulating Factor monkey recombinant produced in *E. coli* is a single, non-glycosylated, polypeptide chain containing 127 amino acids and having a molecular mass of 14.4 kDa. GM-CSF is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

GM-CSF was lyophilized after extensive dialysis against 1x PBS pH 7.4±0.1.

Solubility

It is recommended to reconstitute the lyophilized GMCSF in sterile 18 MΩ-cm H₂O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized GMCSF, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution GMCSF should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity

Greater than 95.0% as determined by (a) Analysis by HPLC, (b) Analysis by SDS-PAGE.

Amino Acid Sequence

APARSPSPGT QPWEHVNAIQ EARRLLNLSR DTAAEMNKTV EVVSEMFDLQ EPSCLQTRLE LYKQGLQGSL TKLKGPLTMM
ASHYKQHCPP TPETSCATQI ITFQSFKENL KDFLLVIPFD CWEPVQE

Activity

The ED50 as determined by the dose-dependant stimulation of the proliferation of human TF-1 cells (human erythroleukemic indicator cell line) is <0.1 ng/ml, corresponding to a specific activity of 10,000,000 IU/mg.



Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.