



## Transforming Growth Factor-Beta 3, His Tag, human recombinant (rHuTGF-b3-His)

**Catalog No:** 97386  
**Lot No:** XXXXX  
**Source:** *Nicotiana benthamiana*  
**Synonyms:** Transforming Growth Factor-beta3, TGFB3, ARVD, FLJ16571, TGF-beta3

### Background

Transforming growth factor betas (TGF Betas) mediate many cell-cell interactions that occur during embryonic development. Three TGF Betas have been identified in mammals. TGF Beta 1, TGF Beta 2 and TGF Beta 3 are each synthesized as precursor proteins that are very similar in that each is cleaved to yield a 112 amino acid polypeptide that remains associated with the latent portion of the molecule.

### Description

TGF-b3 human recombinant produced in plant is a disulfide-linked homodimeric, glycosylated, polypeptide chain containing 118 amino acids and having a molecular mass of 27.2 kDa. TGFB3 is fused to 6xHis Tag at N-terminus and purified by proprietary chromatographic techniques.

### Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

### Formulation

Lyophilized from a concentrated (1 mg/ml) solution containing 50 mM Tris-HCl pH 7.4.

### Solubility

It is recommended to reconstitute the lyophilized TGFB3 in sterile 5 mM HCl and 50 µg/ml BSA at a concentration of 0.05 mg/ml, which can then be further diluted to other aqueous solutions.

### Stability

Lyophilized TGFB3, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution TGFB3 Human should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

### Purity

Greater than 95.0% as determined by SDS-PAGE.

### Amino Acid Sequence

HHHHHALDTN YCFRNLEENC CVRPLYIDFR QDLGKWKVHE PKGYANFCS GPCPYLRSAD TTHSTVLGLY NTLNPEASAS  
PCCVPQDLEP LTILYYVGRT PKVEQLSNMV VKSCKCS

### Activity

The biological activity of TGFB3 is measured in culture by its ability to inhibit the mink lung epithelial (Mv1Lu) cells proliferation. ED50 = 40 ng/ml corresponding to a specific activity of 25,000 units/mg.

## CONTACT US TODAY



### Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.

**CONTACT US TODAY**

BIOMOL GmbH • Kieler Straße 303a • 22525 Hamburg • Germany • [info@biomol.de](mailto:info@biomol.de) • [www.biomol.de](http://www.biomol.de)

Fon: +49 (0)40-853 260 0 • TOLL FREE IN GERMANY: Fon: 0800-246 66 51