



Epithelial Neutrophil-Activating Protein 78, mouse recombinant (CXCL5) (rmCXCL5)

Catalog No: 99959
Lot No: XXXXX
Source: *E. coli*
Synonyms: C-X-C motif chemokine 5, Small-inducible cytokine B5, Cytokine LIX, Cxcl5, Scyb5, LIX, GCP-2, Scyb6, ENA-78, AMCF-II

Background

Chemokine (C-X-C motif) ligand 5 (CXCL5) is a small cytokine belonging to the CXC chemokine family, that is also known as epithelial-derived neutrophil-activating peptide 78 (ENA-78). It is produced following stimulation of cells with the inflammatory cytokines IL-1 or TNF- α . Expression of CXCL5 has also been observed in eosinophils and can be inhibited with the type II interferon IFN- γ . This chemokine stimulates the chemotaxis of neutrophils possessing angiogenic properties. It elicits these effects by interacting with the cell surface chemokine receptor CXCR2. The gene for CXCL5 is encoded on four exons and is located on human chromosome 4 amongst several other CXC chemokine genes. CXCL5 has been implicated in connective tissue remodelling.

Description

Epithelial Neutrophil-Activating Protein 78 Mouse Recombinant, produced in *E. coli*, is a single, non-glycosylated polypeptide chain containing 93 amino acids. It has a molecular mass of 9.8 kDa. The CXCL5 is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered, white, lyophilized (freeze-dried) powder.

Formulation

Lyophilized from a 0.2 μ m filtered concentrated (1.0 mg/ml) solution in 20 mM sodium phosphate buffer (pH 7.4) and 150 mM NaCl.

Solubility

It is recommended to reconstitute the lyophilized ENA-78 in sterile 18 M Ω -cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized ENA-78, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CXCL5 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid Sequence

APSSVIAATE LRCVCLTVTP KINPKLIANL EVIPAGPQCP TVEVIACLKN QKEVCLDPEA PVIKKIIIQK ILGSDKKKAK
RNALAVERTA SVQ

Activity

Determined by its ability to chemoattract human peripheral blood neutrophils, using a concentration range of 10-100 ng/ml, corresponds to a specific activity of 10,000-100,000 IU/mg.

CONTACT US TODAY

BIOMOL GmbH • Kieler Straße 303a • 22525 Hamburg • Germany • info@biomol.de • www.biomol.de

Fon: +49 (0)40-853 260 0 • TOLL FREE IN GERMANY: Fon: 0800-246 66 51



Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.

CONTACT US TODAY

BIOMOL GmbH • Kieler Straße 303a • 22525 Hamburg • Germany • info@biomol.de • www.biomol.de

Fon: +49 (0)40-853 260 0 • TOLL FREE IN GERMANY: Fon: 0800-246 66 51