

Rabbit IgG–Fc Fragment Antibody

Goat Polyclonal
Antigen Affinity Purified
Conjugate DyLight® 488
Catalog No. A120-111D2
Lot No. A120-111D2-7



APPLICATIONS IHC, ICC, F, IF
SPECIES REACTIVITY Rabbit
ISOTYPE IgG
AMOUNT 1 ml at 0.5 mg/ml
STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt
PHYSICAL STATE Liquid
FLUOROPHORE/PROTEIN 6
BUFFER Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide
ORIGIN USA
PRODUCTION PROCEDURES Antiserum was solid phase adsorbed to ensure class specificity. The antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 488.

Antibody concentration was determined by extinction coefficient prior to conjugation: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

By immunoelectrophoresis, this antibody reacts specifically with rabbit IgG. This antibody may cross react with IgG from other species.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry 1:40 – 1:400
Immunocytochemistry 1:40 – 1:500
Flow Cytometry 1:50 – 1:200
Immunofluorescence 1:40 – 1:500


APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

DyLight® 488 is excited at 493 (in PBS) and emits at 518 (in PBS).

DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

ADDITIONAL INFO <https://www.bethyl.com/product/A120-111D2>
Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer Date: December 3, 2018

**Detection of human PCNA by immunofluorescence.**

Sample: FFPE serial sections of human stomach carcinoma.

Primary Antibody: Affinity purified rabbit anti-PCNA (Cat. No. IHC-00012 Lot1) used at a dilution of 1:100 (left) and primary antibody omitted (right). *Secondary Antibody:* Green-fluorescent goat anti-rabbit IgG- Fc Fragment Antibody DyLight® 488 Conjugated (A120-111D2 Lot 3) used at a dilution of 1:50 (10µg/ml). *Counterstain:* DAPI (blue)