Rabbit IgG-Fc Fragment cross-adsorbed Antibody

Goat Polyclonal Conjugate FITC

Antigen Affinity Purified

Catalog No. A120-211F

Lot No. A120-211F-8



APPLICATIONS IHC, ICC, F, IF

SPECIES REACTIVITY Rabbit. Minimum reactivity to human, mouse and rat

AMOUNT 1 ml

CONCENTRATION 0.5 mg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide

FLUOROPHORE/PROTEIN 5.1
ISOTYPE IgG
ORIGIN USA

PRODUCTIONAntiserum was solid phase adsorbed to ensure class specificity. Antiserum was cross adsorbed using human, mouse and rat immunosorbents to remove cross reactive antibodies. The antibody

to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads

and conjugated to fluorescein isothiocyanate (FITC).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

By immunoelectrophoresis, this antibody reacts specifically with rabbit IgG. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to human,

mouse and rat IgG was detected. This antibody may cross react with IgG from other species.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry 1:50 - 1:500

Immunocytochemistry 1:50 – 1:500

Flow Cytometry 1:50 - 1:200

Immunofluorescence 1:50 - 1:500

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

ADDITIONAL INFO https://www.bethyl.com/product/A120-211F

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: September 11, 2019