
AU1 Tag Antibody

AU1 Tag Antibody				
Goat Polyclonal				
Antigen Affinity Purified				
Catalog No.	Catalog No. A190–124A			DETUVI
Lot No.	A190-	124A-1		
APPLICATIONS		WB, IP, ICC, ELISA		
AMOUNT		0.1 ml		
CONCENTRATION		1 mg/ml		
STORAGE/SHELF LIFE		2 - 8° C / 1 year from date of receipt		
PHYSICAL STATE		Liquid		
BUFFER		Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide		
ISOTYPE		IgG		
ORIGIN		USA		
PRODUCTION PROCEDURES		Goats were immunized with AU1 (DTYRYI) conjugated to KLH. Antibody was isolated by affinity chromatography using the peptide immobilized on solid support.		
		Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.		
APPLICATIONS		Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.		
		Western Blot	1:1,000 - 1:30,000	
		Immunoprecipitation	1 – 4 µg/mg lysate	
		Immunocytochemistry	1:100 - 1:400	
		ELISA	1:1,000 - 1:30,000; for coating plates 1:100 -	- 1:500
APPLICATION NOTES		Not all listed applications have been specifically tested by our laboratory.		
ADDITIONAL INFO		Validation by Western Blot was performed using a Western Blot Gel 4–20%. https://www.bethyl.com/product/A190–124A Use the link above to view SDS, a current list of citations, and other product specific information.		

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: September 3, 2019

For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.



Detection of AU1-tagged Protein by western blot.

Samples: 200, 100, or 50 ng of E. coli whole cell lysate expressing a multi-tag fusion protein. Antibodies: Affinity purified, goat anti-AU1 antibody A190-124A used for WB at 0.2 μ g/ml (1:5,000). Detection: Chemiluminescence with an exposure time of 30 seconds.

Bethyl Laboratories, Inc. • 25043 West FM 1097 • Montgomery, TX 77356 • 800.338.9579 • 936.597.6111 • 866.597.6105 (FAX) • www.bethyl.com • technical@bethyl.com