

EWS Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID Q01844

Catalog No. A300-416A


GeneID 2130

Lot No. A300-416A-1



APPLICATIONS	WB				
SPECIES REACTIVITY	Human				
PRESUMED REACTIVITY	Based on 100% sequence identity, this antibody is predicted to react with Mouse				
AMOUNT	100 µl				
CONCENTRATION	1000 µg/ml				
STORAGE/SHELF LIFE	2 - 8° C / 1 year from date of receipt				
PHYSICAL STATE	Liquid				
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide				
ISOTYPE	IgG				
ORIGIN	USA				
PRODUCTION PROCEDURES	<p>Antibody was affinity purified using an epitope specific to EWS immobilized on solid support.</p> <p>The epitope recognized by A300-416A maps to a region between residues 1 and 50 of human Ewing sarcoma breakpoint region 1 using the numbering given in SwissProt entry Q01844 (GeneID 2130).</p> <p>Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.</p>				
APPLICATIONS	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Western Blot</td><td>1:5,000 – 1:15,000</td></tr><tr><td>Immunoprecipitation</td><td>Not recommended. Rabbit anti-EWS Antibodies A300-417A and A300-418A are recommended.</td></tr></table>	Western Blot	1:5,000 – 1:15,000	Immunoprecipitation	Not recommended. Rabbit anti-EWS Antibodies A300-417A and A300-418A are recommended.
Western Blot	1:5,000 – 1:15,000				
Immunoprecipitation	Not recommended. Rabbit anti-EWS Antibodies A300-417A and A300-418A are recommended.				
APPLICATION NOTES	Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.				
ADDITIONAL INFO	<p>https://www.bethyl.com/product/A300-416A</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p>				

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



Detection of human EWS by western blot. *Samples:* Whole cell lysate (50 μ g) from HeLa and HEK293T cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-EWS antibody A300-416A (lot A300-416A-1) used for WB at 0.066 μ g/ml. *Detection:* Chemiluminescence with an exposure time of 3 minutes.