

HDAC2 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP_001518.1

Catalog No. A300-704A


GeneID 3066

Lot No. A300-704A-1



APPLICATIONS	IP
SPECIES REACTIVITY	Human
PRESUMED REACTIVITY	Based on 100% sequence identity, this antibody is predicted to react with Mouse and Chicken
AMOUNT	100 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 - 8° C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	<p>Antibody was affinity purified using an epitope specific to HDAC2 immobilized on solid support.</p> <p>The epitope recognized by A300-704A maps to the N-terminus of human Histone Deacetylase 2 using the numbering given in entry NP_001518.1 (GeneID 3066).</p> <p>Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.</p>
APPLICATIONS	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <p>Western Blot Not recommended. Use rabbit anti-HDAC2 antibody A300-705A.</p> <p>Immunoprecipitation 2 - 5 µg/mg lysate</p>
APPLICATION NOTES	Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).
ADDITIONAL INFO	<p>https://www.bethyl.com/product/A300-704A</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB</p>

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



Detection of human HDAC2 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1 mg for IP, 20% of IP loaded), from HeLa cells. *Antibodies:* Affinity purified rabbit anti-HDAC2 antibody BL2666 (Cat. No. A300-704A) used for IP at 3 μ g/mg lysate. HDAC2 was also immunoprecipitated using rabbit anti-HDAC2 antibody BL2667 (Cat. No. A300-705A) at 3 μ g/mg lysate. For blotting immunoprecipitated HDAC2, BL2667 was used at 1 μ g/ml. *Detection:* Chemiluminescence with an exposure time of 3 seconds.