

KIFC1 /MKLP2 Antibody

Rabbit Polyclonal

Antigen Affinity Purified	RefSeq ID	NP_002254.1
Catalog No. A300-951A	Uniprot ID	Q9BW19
Lot No. 2	GeneID	3833

APPLICATIONS	WB, IP, IHC
SPECIES REACTIVITY	Human
AMOUNT	100 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris–citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to KIFC1 /MKLP2 immobilized on solid support.

The epitope recognized by A300-951A maps to a region between residue 1 and 50 of human Kinesin Family Member C1 (Mitotic Kinesin-Like Protein 2) using the numbering given in entry NP_002254.1 (GeneID 3833).

Immunoglobulin concentration was determined using Beer’s Law where 1 mg/mL IgG has an A280 of 1.4.


APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot	1:2000 – 1:10,000
Immunoprecipitation	6 µg/mg lysate
Immunohistochemistry	1:1000 – 1:5000. Epitope retrieval with Tris–EDTA pH 9.0 is recommended for FFPE tissue sections.

IHC HUMAN CONTROLS Ovarian Carcinoma

ADDITIONAL INFO <https://www.fortislife.com/p/A300-951A>
Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: July 24, 2024




Detection of human KIFC1/MKLP2 by western blot.

Samples: Whole cell lysate (50 µg) from Jurkat, K-562, and CCRF-CEM cells prepared using NETN lysis buffer.


Antibody: Affinity purified rabbit anti-KIFC1/MKLP2 antibody (A300-951A) used for WB at 0.4 µg/ml.

Detection: Chemiluminescence with an exposure time of 30 seconds.



Detection of human KIFC1/MKLP2 by western blot of immunoprecipitates.

Samples: Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from Jurkat cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-KIFC1/MKLP2 antibody (A300-951A) used for IP at 6 µg per reaction. KIFC1-MKLP2 was also immunoprecipitated by a previous lot of this antibody (A300-951A lot 1) and a second antibody against a different epitope of KIFC1/MKLP2 (A300-952A). For blotting immunoprecipitated KIFC1/MKLP2, A300-951A was used at 0.4 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.



Detection of human KIFC1/MKLP2 by immunohistochemistry. *Sample:* FFPE section of human ovarian carcinoma. *Antibody:* Affinity purified rabbit anti-KIFC1/MKLP2 (A300-951A) used at a dilution of 1:5,000 (0.2µg/ml). *Detection:* DAB