SAFB2 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_055464.1

Catalog No. A301-112A GenelD 9667

Lot No. A301-112A-1

APPLICATIONS WB, IP, IHC

SPECIES REACTIVITY Human

AMOUNT 100 μl

CONCENTRATION 200 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to SAFB2 immobilized on solid support.

The epitope recognized by A301-112A maps to a region between residue 175 and 225 of human scaffold attachment factor B2 using the numbering given in entry NP_055464.1 (GeneID 9667).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm

of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000 Immunoprecipitation 2 - 5 µg/mg lysate

Immunohistochemistry 1:200 - 1:1,000. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE tissue sections.

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.

IHC HUMAN CONTROLS Breast Carcinoma, Colon Carcinoma, Non-Small Cell Lung Cancer, Ovarian Carcinoma, Prostate

Carcinoma, Small Cell Lung Cancer

ADDITIONAL INFO https://www.bethvl.com/product/A301-112A

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer


Date: June 21, 2019



SAFB2 Antibody A301-112A



Detection of human SAFB2 by western blot and immunoprecipitation. Samples: Whole cell lysate from HeLa (5, 15 and 50 μ g for WB; 1 mg for IP, 20% of IP loaded) and HEK293T (T: 50 μ g) cells. Antibodies: Affinity purified rabbit anti–SAFB2 antibody A301–112A used for WB at 0.04 μ g/ml (A) and 1 μ g/ml (B) and used for IP at 3 μ g/mg lysate. SAFB2 was also immunoprecipitated by rabbit anti–SAFB2 antibodies A301–113A and BL4609, which recognize downstream epitopes. Detection: Chemiluminescence with exposure times of 3 seconds (A and B).



Detection of human SAFB2 by immunohistochemistry. Sample: FFPE section of human small cell lung cancer. Antibody: Affinity purified rabbit anti- SAFB2 (Cat. No. A301-112A Lot1) used at a dilution of 1:200 (1µg/ml). Detection: DAB